

C_{2h}^2
 $P112_1/m$

No. 11

 $P2_1/m$

 UNIQUE AXIS c
Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3)

General position

 Multiplicity,
 Wyckoff letter,
 Site symmetry

Coordinates

 4 f 1 (1) x, y, z (2) $\bar{x}, \bar{y}, z + \frac{1}{2}$ (3) $\bar{x}, \bar{y}, \bar{z}$ (4) $x, y, \bar{z} + \frac{1}{2}$
I Maximal translationengleiche subgroups

[2] $P11m$ (6)	1; 4		0, 0, 1/4
[2] $P112_1$ (4)	1; 2		
[2] $P\bar{1}$ (2)	1; 3		

II Maximal klassengleiche subgroups

• Enlarged unit cell

[2] $\mathbf{a}' = 2\mathbf{a}$			
$P112_1/a$ (14)	$\langle 3; 2 + (1, 0, 0) \rangle$	$2\mathbf{a}, \mathbf{b}, \mathbf{c}$	
$P112_1/a$ (14)	$\langle 2; 3 + (1, 0, 0) \rangle$	$2\mathbf{a}, \mathbf{b}, \mathbf{c}$	1/2, 0, 0
$P112_1/m$ (11)	$\langle 2; 3 \rangle$	$2\mathbf{a}, \mathbf{b}, \mathbf{c}$	
$P112_1/m$ (11)	$\langle (2; 3) + (1, 0, 0) \rangle$	$2\mathbf{a}, \mathbf{b}, \mathbf{c}$	1/2, 0, 0
[2] $\mathbf{b}' = 2\mathbf{b}$			
$P112_1/b$ (14, $P112_1/a$)	$\langle 3; 2 + (0, 1, 0) \rangle$	$2\mathbf{b}, -\mathbf{a} - 2\mathbf{b}, \mathbf{c}$	
$P112_1/b$ (14, $P112_1/a$)	$\langle 2; 3 + (0, 1, 0) \rangle$	$2\mathbf{b}, -\mathbf{a} - 2\mathbf{b}, \mathbf{c}$	0, 1/2, 0
$P112_1/m$ (11)	$\langle 2; 3 \rangle$	$\mathbf{a}, 2\mathbf{b}, \mathbf{c}$	
$P112_1/m$ (11)	$\langle (2; 3) + (0, 1, 0) \rangle$	$\mathbf{a}, 2\mathbf{b}, \mathbf{c}$	0, 1/2, 0
[2] $\mathbf{a}' = 2\mathbf{a}, \mathbf{b}' = 2\mathbf{b}$			
$C112_1/e$ (14, $P112_1/a$)	$\langle 3; 2 + (1, 0, 0) \rangle$	$2\mathbf{a}, -\mathbf{a} + \mathbf{b}, \mathbf{c}$	
$C112_1/e$ (14, $P112_1/a$)	$\langle 2; 3 + (1, 0, 0) \rangle$	$2\mathbf{a}, -\mathbf{a} + \mathbf{b}, \mathbf{c}$	1/2, 0, 0
$C112_1/m$ (11, $P112_1/m$)	$\langle 2; 3 \rangle$	$2\mathbf{a}, -\mathbf{a} + \mathbf{b}, \mathbf{c}$	
$C112_1/m$ (11, $P112_1/m$)	$\langle (2; 3) + (1, 0, 0) \rangle$	$2\mathbf{a}, -\mathbf{a} + \mathbf{b}, \mathbf{c}$	1/2, 0, 0
[3] $\mathbf{c}' = 3\mathbf{c}$			
$P112_1/m$ (11)	$\langle 3; 2 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	
$P112_1/m$ (11)	$\langle 2 + (0, 0, 1); 3 + (0, 0, 2) \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	0, 0, 1
$P112_1/m$ (11)	$\langle 2 + (0, 0, 1); 3 + (0, 0, 4) \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	0, 0, 2
[3] $\mathbf{a}' = 3\mathbf{a}$			
$P112_1/m$ (11)	$\langle 2; 3 \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	
$P112_1/m$ (11)	$\langle (2; 3) + (2, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	1, 0, 0
$P112_1/m$ (11)	$\langle (2; 3) + (4, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	2, 0, 0
[3] $\mathbf{a}' = 3\mathbf{a}, \mathbf{b}' = -\mathbf{a} + \mathbf{b}$			
$P112_1/m$ (11)	$\langle 2; 3 \rangle$	$3\mathbf{a}, -\mathbf{a} + \mathbf{b}, \mathbf{c}$	
$P112_1/m$ (11)	$\langle (2; 3) + (2, 0, 0) \rangle$	$3\mathbf{a}, -\mathbf{a} + \mathbf{b}, \mathbf{c}$	1, 0, 0
$P112_1/m$ (11)	$\langle (2; 3) + (4, 0, 0) \rangle$	$3\mathbf{a}, -\mathbf{a} + \mathbf{b}, \mathbf{c}$	2, 0, 0
[3] $\mathbf{a}' = 3\mathbf{a}, \mathbf{b}' = -2\mathbf{a} + \mathbf{b}$			
$P112_1/m$ (11)	$\langle 2; 3 \rangle$	$3\mathbf{a}, -2\mathbf{a} + \mathbf{b}, \mathbf{c}$	
$P112_1/m$ (11)	$\langle (2; 3) + (2, 0, 0) \rangle$	$3\mathbf{a}, -2\mathbf{a} + \mathbf{b}, \mathbf{c}$	1, 0, 0
$P112_1/m$ (11)	$\langle (2; 3) + (4, 0, 0) \rangle$	$3\mathbf{a}, -2\mathbf{a} + \mathbf{b}, \mathbf{c}$	2, 0, 0
[3] $\mathbf{b}' = 3\mathbf{b}$			
$P112_1/m$ (11)	$\langle 2; 3 \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	
$P112_1/m$ (11)	$\langle (2; 3) + (0, 2, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 1, 0
$P112_1/m$ (11)	$\langle (2; 3) + (0, 4, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 2, 0

• Series of maximal isomorphic subgroups

[p] $\mathbf{c}' = p\mathbf{c}$			
$P112_1/m$ (11)	$\langle 2 + (0, 0, \frac{p}{2} - \frac{1}{2}); 3 + (0, 0, 2u) \rangle$	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	0, 0, u
	$p > 2; 0 \leq u < p$		
	p conjugate subgroups for the prime p		
[p] $\mathbf{a}' = p\mathbf{a}, \mathbf{b}' = -q\mathbf{a} + \mathbf{b}$			
$P112_1/m$ (11)	$\langle (2; 3) + (2u, 0, 0) \rangle$	$p\mathbf{a}, -q\mathbf{a} + \mathbf{b}, \mathbf{c}$	$u, 0, 0$
	$p > 2; 0 \leq q < p; 0 \leq u < p$		
	p conjugate subgroups for each pair of q and prime p		
[p] $\mathbf{b}' = p\mathbf{b}$			
$P112_1/m$ (11)	$\langle (2; 3) + (0, 2u, 0) \rangle$	$\mathbf{a}, p\mathbf{b}, \mathbf{c}$	0, $u, 0$
	$p > 2; 0 \leq u < p$		
	p conjugate subgroups for the prime p		

(Continued on the facing page)

I Minimal translationengleiche supergroups

[2] $Pmma$ (51); [2] $Pbcm$ (57); [2] $Pmnn$ (59); [2] $Pnma$ (62); [2] $Cmcm$ (63); [3] $P6_3/m$ (176)

II Minimal non-isomorphic klassengleiche supergroups**• Additional centring translations**

[2] $C12/m1$ (12); [2] $A12/m1$ (12, $C12/m1$); [2] $I12/m1$ (12, $C12/m1$)

• Decreased unit cell

[2] $\mathbf{b}' = \frac{1}{2}\mathbf{b}$ $P12/m1$ (10)

I Minimal translationengleiche supergroups

[2] $Pmma$ (51); [2] $Pbcm$ (57); [2] $Pmnn$ (59); [2] $Pnma$ (62); [2] $Cmcm$ (63); [3] $P6_3/m$ (176)

II Minimal non-isomorphic klassengleiche supergroups**• Additional centring translations**

[2] $A112/m$ (12); [2] $B112/m$ (12, $A112/m$); [2] $I112/m$ (12, $A112/m$)

• Decreased unit cell

[2] $\mathbf{c}' = \frac{1}{2}\mathbf{c}$ $P112/m$ (10)