

$P4/nbm$ 

No. 125

 $P4/n2/b2/m$  $D_{4h}^3$ ORIGIN CHOICE 1, Origin at 422 at  $4/n22/g$ , at  $-\frac{1}{4}, -\frac{1}{4}, 0$  from centre ( $2/m$ )**Generators selected** (1);  $t(1,0,0)$ ;  $t(0,1,0)$ ;  $t(0,0,1)$ ; (2); (3); (5); (9)**General position**Multiplicity,  
Wyckoff letter,  
Site symmetry

Coordinates

16	$n$	1	(1) $x, y, z$	(2) $\bar{x}, \bar{y}, z$	(3) $\bar{y}, x, z$	(4) $y, \bar{x}, z$
			(5) $\bar{x}, y, \bar{z}$	(6) $x, \bar{y}, \bar{z}$	(7) $y, x, \bar{z}$	(8) $\bar{y}, \bar{x}, \bar{z}$
			(9) $\bar{x} + \frac{1}{2}, \bar{y} + \frac{1}{2}, \bar{z}$	(10) $x + \frac{1}{2}, y + \frac{1}{2}, \bar{z}$	(11) $y + \frac{1}{2}, \bar{x} + \frac{1}{2}, \bar{z}$	(12) $\bar{y} + \frac{1}{2}, x + \frac{1}{2}, \bar{z}$
			(13) $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, z$	(14) $\bar{x} + \frac{1}{2}, y + \frac{1}{2}, z$	(15) $\bar{y} + \frac{1}{2}, \bar{x} + \frac{1}{2}, z$	(16) $y + \frac{1}{2}, x + \frac{1}{2}, z$

**I Maximal translationengleiche subgroups**

[2] $P\bar{4}b2$ (117)	1; 2; 7; 8; 11; 12; 13; 14	0, 1/2, 0
[2] $P\bar{4}2m$ (111)	1; 2; 5; 6; 11; 12; 15; 16	0, 1/2, 0
[2] $P4bm$ (100)	1; 2; 3; 4; 13; 14; 15; 16	
[2] $P422$ (89)	1; 2; 3; 4; 5; 6; 7; 8	
[2] $P4/n11$ (85, $P4/n$ )	1; 2; 3; 4; 9; 10; 11; 12	0, 1/2, 0
[2] $P2/n12/m$ (67, $Cmme$ )	1; 2; 7; 8; 9; 10; 15; 16	$\mathbf{a} - \mathbf{b}, \mathbf{a} + \mathbf{b}, \mathbf{c}$
[2] $P2/n2/b1$ (50, $Pban$ )	1; 2; 5; 6; 9; 10; 13; 14	1/4, 3/4, 0

**II Maximal klassengleiche subgroups****• Enlarged unit cell**[2]  $\mathbf{c}' = 2\mathbf{c}$ 

$P4_2/nnm$ (134)	$\langle 2; 9; (3; 5) + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	0, 1/2, 1/2
$P4_2/nnm$ (134)	$\langle 2; 5; (3; 9) + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	0, 1/2, 0
$P4_2/nbc$ (133)	$\langle 2; 5; 9; 3 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	0, 1/2, 1/2
$P4_2/nbc$ (133)	$\langle 2; (3; 5; 9) + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	0, 1/2, 0
$P4/nnc$ (126)	$\langle 2; 3; 9; 5 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	0, 0, 1/2
$P4/nnc$ (126)	$\langle 2; 3; 5; 9 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	
$P4/nbm$ (125)	$\langle 2; 3; 5; 9 \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	
$P4/nbm$ (125)	$\langle 2; 3; (5; 9) + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	0, 0, 1/2
[3] $\mathbf{c}' = 3\mathbf{c}$	$\left\{ \begin{array}{l} P4/nbm (125) \\ P4/nbm (125) \\ P4/nbm (125) \end{array} \right.$	$\langle 2; 3; 5; 9 \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$
		$\langle 2; 3; (5; 9) + (0, 0, 2) \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$
		$\langle 2; 3; (5; 9) + (0, 0, 4) \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$

**• Series of maximal isomorphic subgroups**[p]  $\mathbf{c}' = p\mathbf{c}$ 

$P4/nbm$ (125)	$\langle 2; 3; (5; 9) + (0, 0, 2u) \rangle$	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	0, 0, u
	$p > 2; 0 \leq u < p$		

*p conjugate subgroups for the prime p*

[ $p^2$ ]  $\mathbf{a}' = p\mathbf{a}$ ,  $\mathbf{b}' = p\mathbf{b}$ 

$P4/nbm$ (125)	$\langle 2 + (2u, 2v, 0); 3 + (u + v, -u + v, 0); 5 + (2u, 0, 0); 9 + (\frac{p}{2} - \frac{1}{2} + 2u, \frac{p}{2} - \frac{1}{2} + 2v, 0) \rangle$	$p\mathbf{a}, p\mathbf{b}, \mathbf{c}$	$u, v, 0$
	$p > 2; 0 \leq u < p; 0 \leq v < p$		
	$p^2$ conjugate subgroups for the prime p		

**I Minimal translationengleiche supergroups**

none

**II Minimal non-isomorphic klassengleiche supergroups****• Additional centring translations**[2]  $C4/mmm$  (123,  $P4/mmm$ ); [2]  $I4/mcm$  (140)**• Decreased unit cell**

none