

$I\bar{4}_1/amd$

No. 141

 $I\bar{4}_1/a2/m2/d$ D_{4h}^{19} ORIGIN CHOICE 1, Origin at $\bar{4}m2$, at $0, \frac{1}{4}, -\frac{1}{8}$ from centre ($2/m$)**Generators selected** (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; $t(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$; (2); (3); (5); (9)**General position**Multiplicity,
Wyckoff letter,
Site symmetry**Coordinates**

$(0,0,0) + (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}) +$

32	i	1	(1) x, y, z	(2) $\bar{x} + \frac{1}{2}, \bar{y} + \frac{1}{2}, z + \frac{1}{2}$	(3) $\bar{y}, x + \frac{1}{2}, z + \frac{1}{4}$	(4) $y + \frac{1}{2}, \bar{x}, z + \frac{3}{4}$
			(5) $\bar{x} + \frac{1}{2}, y, \bar{z} + \frac{3}{4}$	(6) $x, \bar{y} + \frac{1}{2}, \bar{z} + \frac{1}{4}$	(7) $y + \frac{1}{2}, x + \frac{1}{2}, \bar{z} + \frac{1}{2}$	(8) $\bar{y}, \bar{x}, \bar{z}$
			(9) $\bar{x}, \bar{y} + \frac{1}{2}, \bar{z} + \frac{1}{4}$	(10) $x + \frac{1}{2}, y, \bar{z} + \frac{3}{4}$	(11) y, \bar{x}, \bar{z}	(12) $\bar{y} + \frac{1}{2}, x + \frac{1}{2}, \bar{z} + \frac{1}{2}$
			(13) $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, z + \frac{1}{2}$	(14) \bar{x}, y, z	(15) $\bar{y} + \frac{1}{2}, \bar{x}, z + \frac{3}{4}$	(16) $y, x + \frac{1}{2}, z + \frac{1}{4}$

I Maximal translationengleiche subgroups

[2] $I\bar{4}2d$ (122)	(1; 2; 5; 6; 11; 12; 15; 16) +			
[2] $I\bar{4}m2$ (119)	(1; 2; 7; 8; 11; 12; 13; 14) +			
[2] $I\bar{4}_1md$ (109)	(1; 2; 3; 4; 13; 14; 15; 16) +			
[2] $I\bar{4}_122$ (98)	(1; 2; 3; 4; 5; 6; 7; 8) +			
[2] $I\bar{4}_1/a11$ (88, $I\bar{4}_1/a$)	(1; 2; 3; 4; 9; 10; 11; 12) +			
[2] $I2/a2/m1$ (74, $Imma$)	(1; 2; 5; 6; 9; 10; 13; 14) +			0, 1/4, 1/8
[2] $I2/a12/d$ (70, $Fddd$)	(1; 2; 7; 8; 9; 10; 15; 16) +			0, 1/2, 1/4
		a - b, a + b, c		

II Maximal klassengleiche subgroups**• Loss of centring translations**

none

• Enlarged unit cell

[3] $c' = 3c$				
$\left\{ \begin{array}{l} I\bar{4}_1/amd \text{ (141)} \\ I\bar{4}_1/amd \text{ (141)} \\ I\bar{4}_1/amd \text{ (141)} \end{array} \right.$	$\langle (2; 9) + (1, 0, 1); 3 + (\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}); 5 + (1, 0, 2) \rangle$ $\langle 2 + (1, 0, 1); 3 + (\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}); 5 + (1, 0, 4); 9 + (1, 0, 3) \rangle$ $\langle 2 + (1, 0, 1); 3 + (\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}); 5 + (1, 0, 6); 9 + (1, 0, 5) \rangle$	a, b, 3c	1/2, 0, 1/4	
		a, b, 3c	1/2, 0, 5/4	
		a, b, 3c	1/2, 0, 9/4	

• Series of maximal isomorphic subgroups**[p] $c' = pc$**

$I\bar{4}_1/amd$ (141)	$\langle 2 + (1, 0, \frac{p}{2} - \frac{1}{2}); 3 + (\frac{1}{2}, -\frac{1}{2}, \frac{p}{4} - \frac{1}{4}); 5 + (1, 0, \frac{3p}{4} - \frac{1}{4} + 2u); 9 + (1, 0, \frac{p}{4} + \frac{1}{4} + 2u) \rangle$ $p > 2; 0 \leq u < p$ p conjugate subgroups for prime $p \equiv 3 \pmod{4}$	a, b, pc	1/2, 0, 1/4 + u
$I\bar{4}_1/amd$ (141)	$\langle 2 + (0, 0, \frac{p}{2} - \frac{1}{2}); 3 + (0, 0, \frac{p}{4} - \frac{1}{4}); 5 + (0, 0, \frac{3p}{4} - \frac{3}{4} + 2u); 9 + (0, 0, \frac{p}{4} - \frac{1}{4} + 2u) \rangle$ $p > 4; 0 \leq u < p$ p conjugate subgroups for prime $p \equiv 1 \pmod{4}$	a, b, pc	0, 0, u
$[p^2]$ $\mathbf{a}' = p\mathbf{a}$, $\mathbf{b}' = p\mathbf{b}$			
$I\bar{4}_1/amd$ (141)	$\langle 2 + (\frac{p}{2} - \frac{1}{2} + 2u, \frac{p}{2} - \frac{1}{2} + 2v, 0); 3 + (u + v, \frac{p}{2} - \frac{1}{2} - u + v, 0); 5 + (\frac{p}{2} - \frac{1}{2} + 2u, 0, 0); 9 + (2u, \frac{p}{2} - \frac{1}{2} + 2v, 0) \rangle$ $p > 2; 0 \leq u < p; 0 \leq v < p$ p^2 conjugate subgroups for the prime p	$p\mathbf{a}, p\mathbf{b}, \mathbf{c}$	$u, v, 0$

I Minimal translationengleiche supergroups**[3] $Fd\bar{3}m$ (227)****II Minimal non-isomorphic klassengleiche supergroups****• Additional centring translations**

none

• Decreased unit cell**[2] $c' = \frac{1}{2}\mathbf{c}$ $C\bar{4}_2/emd$ (134, $P\bar{4}_2/nm$)**