

## 10. POINT GROUPS AND CRYSTAL CLASSES





Fig. 10.1.3.2. Maximal subgroups and minimal supergroups of the three-dimensional crystallographic point groups. Solid lines indicate maximal normal subgroups; double or triple solid lines mean that there are two or three maximal normal subgroups with the same symbol. Dashed lines refer to sets of maximal conjugate subgroups. The group orders are given on the left. Full Hermann-Mauguin symbols are used.

symbols' 222. and 2.22.



(2) Similarly, group 432 has one maximal normal subgroup, 23.

Figs. 10.1.3.1 and 10.1.3.2 show that there exist two 'summits' in both two and three dimensions from which all other point groups can be derived by 'chains' of maximal subgroups. These summits are formed by the square and the hexagonal holohedry in two dimensions and by the cubic and the hexagonal holohedry in three dimensions.

The sub- and supergroups of the point groups are useful both in their own right and as basis of the *translationengleiche* or *t* subgroups and supergroups of space groups; this is set out in Section 2.2.15. Tables of the sub- and supergroups of the plane groups and space groups are contained in Parts 6 and 7. A general discussion of sub- and supergroups of crystallographic groups, together with further explanations and examples, is given in Section 8.3.3.

### 10.1.4. Noncrystallographic point groups

#### 10.1.4.1. Description of general point groups

In Sections 10.1.2 and 10.1.3, only the 32 *crystallographic* point groups (crystal classes) are considered. In addition, infinitely many *noncrystallographic* point groups exist that are of interest as possible symmetries of molecules and of quasicrystals and as

approximate local site symmetries in crystals. Crystallographic and noncrystallographic point groups are collected here under the name *general point groups*. They are reviewed in this section and listed in Tables 10.1.4.1 to 10.1.4.3.

Because of the infinite number of these groups only *classes of general point groups* (*general classes*)\* can be listed. They are grouped into *general systems*, which are similar to the crystal systems. The 'general classes' are of two kinds: in the cubic, icosahedral, circular, cylindrical and spherical system, each general class contains *one* point group only, whereas in the  $4N$ -gonal,  $(2N + 1)$ -gonal and  $(4N + 2)$ -gonal system, each general class contains *infinitely* many point groups, which differ in their principal  $n$ -fold symmetry axis, with  $n = 4, 8, 12, \dots$  for the  $4N$ -gonal system,  $n = 1, 3, 5, \dots$  for the  $(2N + 1)$ -gonal system and  $n = 2, 6, 10, \dots$  for the  $(4N + 2)$ -gonal system.

Furthermore, some general point groups are of order infinity because they contain symmetry axes (rotation or rotoinversion axes) of order infinity† ( $\infty$ -fold axes). These point groups occur in the

\* The 'classes of general point groups' are not the same as the commonly used 'crystal classes' because some of them contain point groups of *different orders*. All these orders, however, follow a common scheme. In this sense, the 'general classes' are an extension of the concept of (geometric) crystal classes. For example, the general class *mmm* of the  $4N$ -gonal system contains the point groups *4mm* (tetragonal), *8mm* (octagonal), *12mm* (dodecagonal), *16mm* etc.

† The axes of order infinity, as considered here, do not correspond to cyclic groups (as do the axes of finite order) because there is no smallest rotation from which all other rotations can be derived as higher powers, *i.e.* by successive application. Instead, rotations of all possible angles exist. Nevertheless, it is customary to symbolize these axes as  $\infty$  or  $C_\infty$ ; note that the Hermann-Mauguin symbols  $\infty/m$  and  $\infty$  are equivalent, and so are the Schoenflies symbols  $C_{\infty h}$ ,  $S_\infty$  and  $C_{\infty i}$ . (There exist also axes of order infinity that do correspond to cyclic groups, namely axes based upon smallest rotations with irrational values of the rotation angle.)

## 10.1. CRYSTALLOGRAPHIC AND NONCRYSTALLOGRAPHIC POINT GROUPS

circular system (two dimensions) and in the cylindrical and spherical systems (three dimensions).

The Hermann–Mauguin and Schoenflies symbols for the general point groups follow the rules of the crystallographic point groups (*cf.* Section 2.2.4 and Chapter 12.1). This extends also to the infinite groups where symbols like  $\infty m$  or  $C_{\infty v}$  are immediately obvious.

In *two dimensions* (Table 10.1.4.1), the eight general classes are collected into three systems. Two of these, the  $4N$ -gonal and the  $(4N + 2)$ -gonal systems, contain only point groups of finite order with one  $n$ -fold rotation point each. These systems are generalizations of the square and hexagonal crystal systems. The circular system consists of two infinite point groups, with one  $\infty$ -fold rotation point each.

In *three dimensions* (Table 10.1.4.2), the 28 general classes are collected into seven systems. Three of these, the  $4N$ -gonal, the  $(2N + 1)$ -gonal and the  $(4N + 2)$ -gonal systems,\* contain only point groups of finite order with one principal  $n$ -fold symmetry axis each. These systems are generalizations of the tetragonal, trigonal, and hexagonal crystal systems (*cf.* Table 10.1.1.2). The five cubic groups are well known as crystallographic groups. The two icosahedral groups of orders 60 and 120, characterized by special combinations of twofold, threefold and fivefold symmetry axes, are discussed in more detail below. The groups of the cylindrical and the spherical systems are all of order infinity; they describe the symmetries of cylinders, cones, rotation ellipsoids, spheres *etc.*

It is possible to define the three-dimensional point groups on the basis of either rotoinversion axes  $\bar{n}$  or rotoreflection axes  $\tilde{n}$ . The equivalence between these two descriptions is apparent from the following examples:

$$\begin{array}{l} n = 4N \quad : \quad \bar{4} = \tilde{4} \quad \bar{8} = \tilde{8} \quad \dots \quad \bar{n} = \tilde{n} \\ n = 2N + 1 \quad : \quad \bar{1} = \tilde{2} \quad \bar{3} = \tilde{6} = 3 \times \bar{1} \dots \quad \bar{n} = \tilde{2n} = n \times \bar{1} \\ n = 4N + 2 \quad : \quad \bar{2} = \tilde{1} = m \quad \bar{6} = \tilde{3} = 3/m \quad \dots \quad \bar{n} = \tilde{\frac{1}{2}n} = \frac{1}{2}n/m. \end{array}$$

In the present tables, the standard convention of using rotoinversion axes is followed.

Tables 10.1.4.1 and 10.1.4.2 contain for each class its general Hermann–Mauguin and Schoenflies symbols, the group order and the names of the general face form and its dual, the general point form.† Special and limiting forms are not given, nor are ‘Miller indices’ ( $hkl$ ) and point coordinates  $x, y, z$ . They can be derived easily from Tables 10.1.2.1 and 10.1.2.2 for the crystallographic groups.‡

### 10.1.4.2. The two icosahedral groups

The two point groups 235 and  $m\bar{3}5$  of the icosahedral system (orders 60 and 120) are of particular interest among the noncrystallographic groups because of the occurrence of fivefold axes and their increasing importance as symmetries of molecules (viruses), of quasicrystals, and as approximate local site symmetries in crystals (alloys,  $B_{12}$  icosahedron). Furthermore, they contain as special forms the two noncrystallographic *platonic solids*, the

regular icosahedron (20 faces, 12 vertices) and its dual, the regular pentagon-dodecahedron (12 faces, 20 vertices).

The icosahedral groups (*cf.* diagrams in Table 10.1.4.3) are characterized by six fivefold axes that include angles of  $63.43^\circ$ . Each fivefold axis is surrounded by five threefold and five twofold axes, with angular distances of  $37.38^\circ$  between a fivefold and a threefold axis and of  $31.72^\circ$  between a fivefold and a twofold axis. The angles between neighbouring threefold axes are  $41.81^\circ$ , between neighbouring twofold axes  $36^\circ$ . The smallest angle between a threefold and a twofold axis is  $20.90^\circ$ .

Each of the six fivefold axes is perpendicular to five twofold axes; there are thus six maximal conjugate pentagonal subgroups of types 52 (for 235) and  $\bar{5}m$  (for  $m\bar{3}5$ ) with index [6]. Each of the ten threefold axes is perpendicular to three twofold axes, leading to ten maximal conjugate trigonal subgroups of types 32 (for 235) and  $\bar{3}m$  (for  $m\bar{3}5$ ) with index [10]. There occur, furthermore, five maximal conjugate cubic subgroups of types 23 (for 235) and  $m\bar{3}$  (for  $m\bar{3}5$ ) with index [5].

The two icosahedral groups are listed in Table 10.1.4.3, in a form similar to the cubic point groups in Table 10.1.2.2. Each group is illustrated by stereographic projections of the symmetry elements and the general face poles (general points); the complete sets of symmetry elements are listed below the stereograms. Both groups are referred to a cubic coordinate system, with the coordinate axes along three twofold rotation axes and with four threefold axes along the body diagonals. This relation is well brought out by symbolizing these groups as 235 and  $m\bar{3}5$  instead of the customary symbols 532 and  $\bar{5}3m$ .

The table contains also the multiplicities, the Wyckoff letters and the names of the general and special face forms and their duals, the point forms, as well as the oriented face- and site-symmetry symbols. In the icosahedral ‘holohedry’  $m\bar{3}5$ , the *special* ‘Wyckoff position’  $60d$  occurs in three realizations, *i.e.* with three types of polyhedra. In 235, however, these three types of polyhedra are different realizations of the limiting *general* forms, which depend on the location of the poles with respect to the axes 2, 3 and 5. For this reason, the three entries are connected by braces; *cf.* Section 10.1.2.4, *Notes on crystal and point forms*, item (viii).

Not included are the sets of equivalent Miller indices and point coordinates. Instead, only the ‘initial’ triplets ( $hkl$ ) and  $x, y, z$  for each type of form are listed. The complete sets of indices and coordinates can be obtained in two steps§ as follows:

(i) For the face forms the cubic point groups 23 and  $m\bar{3}$  (Table 10.1.2.2), and for the point forms the cubic space groups  $P23$  (195) and  $Pm\bar{3}$  (200) have to be considered. For each ‘initial’ triplet ( $hkl$ ), the set of Miller indices of the (general or special) crystal form with the same face symmetry in 23 (for group 235) or  $m\bar{3}$  (for  $m\bar{3}5$ ) is taken. For each ‘initial’ triplet  $x, y, z$ , the coordinate triplets of the (general or special) position with the same site symmetry in  $P23$  or  $Pm\bar{3}$  are taken; this procedure is similar to that described in Section 10.1.2.3 for the crystallographic point forms.

(ii) To obtain the complete set of icosahedral Miller indices and point coordinates, the ‘cubic’ ( $hkl$ ) triplets (as rows) and  $x, y, z$  triplets (as columns) have to be multiplied with the identity matrix and with

(a) the matrices  $Y, Y^2, Y^3$  and  $Y^4$  for the Miller indices;

\* Here, the  $(2N + 1)$ -gonal and the  $(4N + 2)$ -gonal systems are distinguished in order to bring out the analogy with the trigonal and the hexagonal crystal systems. They could equally well be combined into one, in correspondence with the hexagonal ‘crystal family’ (*cf.* Chapter 2.1).

† The noncrystallographic face and point forms are extensions of the corresponding crystallographic forms: *cf.* Section 10.1.2.4, *Notes on crystal and point forms*. The name *streptohedron* applies to the general face forms of point groups  $\bar{n}$  with  $n = 4N$  and  $n = 2N + 1$ ; it is thus a generalization of the tetragonal disphenoid or tetragonal tetrahedron (4) and the rhombohedron ( $\bar{3}$ ).

‡ The term ‘Miller indices’ is used here also for the noncrystallographic point groups. Note that these indices do not have to be integers or rational numbers, as for the crystallographic point groups. Irrational indices, however, can always be closely approximated by integers, quite often even by small integers.

§ A one-step procedure applies to the icosahedral ‘Wyckoff position’  $12a$ , the face poles and points of which are located on the fivefold axes. Here, step (ii) is redundant and can be omitted. The forms  $\{01\tau\}$  and  $0, y, \tau y$  are contained in the cubic point groups 23 and  $m\bar{3}$  and in the cubic space groups  $P23$  and  $Pm\bar{3}$  as limiting cases of Wyckoff positions  $\{0kl\}$  and  $0, y, z$  with specialized (irrational) values of the indices and coordinates. In geometric terms, the regular pentagon-dodecahedron is a noncrystallographic ‘limiting polyhedron’ of the ‘crystallographic’ pentagon-dodecahedron and the regular icosahedron is a ‘limiting polyhedron’ of the ‘irregular’ icosahedron (*cf.* Section 10.1.2.2, *Crystal and point forms*).

10. POINT GROUPS AND CRYSTAL CLASSES

Table 10.1.4.1. *Classes of general point groups in two dimensions ( $N = \text{integer} \geq 0$ )*

General Hermann–Mauguin symbol	Order of group	General edge form	General point form	Crystallographic groups
4 <i>N</i> -gonal system ( <i>n</i> -fold rotation point with $n = 4N$ )				
<i>n</i> <i>nmm</i>	<i>n</i> $2n$	Regular <i>n</i> -gon Semiregular di- <i>n</i> -gon	Regular <i>n</i> -gon Truncated <i>n</i> -gon	4 <i>4mm</i>
$(4N + 2)$ -gonal system ( <i>n</i> -fold or $\frac{1}{2}n$ -fold rotation point with $n = 4N + 2$ )				
$\frac{1}{2}n$ $\frac{1}{2}nm$ <i>n</i> <i>nmm</i>	$\frac{1}{2}n$ <i>n</i> <i>n</i> $2n$	Regular $\frac{1}{2}n$ -gon Semiregular di- $\frac{1}{2}n$ -gon Regular <i>n</i> -gon Semiregular di- <i>n</i> -gon	Regular $\frac{1}{2}n$ -gon Truncated $\frac{1}{2}n$ -gon Regular <i>n</i> -gon Truncated <i>n</i> -gon	1, 3 <i>m</i> , $3m$ 2, 6 $2mm$ , $6mm$
Circular system *				
$\infty$ $\infty m$	$\infty$ $\infty$	Rotating circle Stationary circle	Rotating circle Stationary circle	– –

\* A rotating circle has no mirror lines; there exist two enantiomorphic circles with opposite senses of rotation. A stationary circle has infinitely many mirror lines through its centre.

Table 10.1.4.2. *Classes of general point groups in three dimensions ( $N = \text{integer} \geq 0$ )*

Short general Hermann–Mauguin symbol, followed by full symbol where different	Schoenflies symbol	Order of group	General face form	General point form	Crystallographic groups
$4N$ -gonal system (single <i>n</i> -fold symmetry axis with $n = 4N$ )					
<i>n</i> $\bar{n}$ <i>n/m</i> <i>n22</i> <i>nmm</i> $\bar{n}2m$ <i>n/mmm</i> , $\frac{n\ 2\ 2}{m\ m\ m}$	$C_n$ $S_n$ $C_{nh}$ $D_n$ $C_{nv}$ $D_{\frac{1}{2}nd}$ $D_{nh}$	<i>n</i> <i>n</i> $2n$ $2n$ $2n$ $2n$ $4n$	<i>n</i> -gonal pyramid $\frac{1}{2}n$ -gonal streptohedron <i>n</i> -gonal dipyramid <i>n</i> -gonal trapezohedron Di- <i>n</i> -gonal pyramid <i>n</i> -gonal scalenohedron Di- <i>n</i> -gonal dipyramid	Regular <i>n</i> -gon $\frac{1}{2}n$ -gonal antiprism <i>n</i> -gonal prism Twisted <i>n</i> -gonal antiprism Truncated <i>n</i> -gon $\frac{1}{2}n$ -gonal antiprism sliced off by pinacoid Edge-truncated <i>n</i> -gonal prism	4 $\frac{4}{4}$ $4/m$ 422 $4mm$ $\bar{4}2m$ $4/mmm$
$(2N + 1)$ -gonal system (single <i>n</i> -fold symmetry axis with $n = 2N + 1$ )					
<i>n</i> $\bar{n} = n \times \bar{1}$ <i>n2</i> <i>nm</i> $\bar{n}m$ , $\bar{n}\frac{2}{m}$	$C_n$ $C_{ni}$ $D_n$ $C_{nv}$ $D_{nd}$	<i>n</i> $2n$ $2n$ $2n$ $4n$	<i>n</i> -gonal pyramid <i>n</i> -gonal streptohedron <i>n</i> -gonal trapezohedron Di- <i>n</i> -gonal pyramid Di- <i>n</i> -gonal scalenohedron	Regular <i>n</i> -gon <i>n</i> -gonal antiprism Twisted <i>n</i> -gonal antiprism Truncated <i>n</i> -gon <i>n</i> -gonal antiprism sliced off by pinacoid	1, 3 $\bar{1}$ , $\bar{3} = 3 \times \bar{1}$ 32 $3m$ $\bar{3}m$
$(4N + 2)$ -gonal system (single <i>n</i> -fold symmetry axis with $n = 4N + 2$ )					
<i>n</i> $\bar{n} = \frac{1}{2}n/m$ <i>n/m</i> <i>n22</i> <i>nmm</i> $\bar{n}2m = \frac{1}{2}n/m2m$ <i>n/mmm</i> , $\frac{n\ 2\ 2}{m\ m\ m}$	$C_n$ $C_{\frac{1}{2}nh}$ $C_{nh}$ $D_n$ $C_{nv}$ $D_{\frac{1}{2}nh}$ $D_{nh}$	<i>n</i> <i>n</i> $2n$ $2n$ $2n$ $2n$ $4n$	<i>n</i> -gonal pyramid $\frac{1}{2}n$ -gonal dipyramid <i>n</i> -gonal dipyramid <i>n</i> -gonal trapezohedron Di- <i>n</i> -gonal pyramid Di- $\frac{1}{2}n$ -gonal dipyramid Di- <i>n</i> -gonal dipyramid	Regular <i>n</i> -gon $\frac{1}{2}n$ -gonal prism <i>n</i> -gonal prism Twisted <i>n</i> -gonal antiprism Truncated <i>n</i> -gon Truncated $\frac{1}{2}n$ -gonal prism Edge-truncated <i>n</i> -gonal prism	2, 6 $\bar{2} \equiv m$ , $\bar{6} \equiv 3/m$ $2/m$ , $6/m$ 222, 622 $mm2$ , $6mm$ $\bar{6}2m$ $mmm$ , $6/mmm$

## 10.1. CRYSTALLOGRAPHIC AND NONCRYSTALLOGRAPHIC POINT GROUPS

Table 10.1.4.2. *Classes of general point groups in three dimensions ( $N = \text{integer} \geq 0$ ) (cont.)*

Short general Hermann–Mauguin symbol, followed by full symbol where different	Schoenflies symbol	Order of group	General face form	General point form	Crystallographic groups
Cubic system (for details see Table 10.1.2.2)					
$23$ $m\bar{3}, \frac{2}{m}\bar{3}$	$T$ $T_h$	12 24	Pentagon-tritetrahedron Didodecahedron	Snub tetrahedron Cube & octahedron & pentagon-dodecahedron	23 $m\bar{3}$
432 $\bar{4}3m$	$O$ $T_d$	24 24	Pentagon-trioctahedron Hexatetrahedron	Snub cube Cube truncated by two tetrahedra	432 $\bar{4}3m$
$m\bar{3}m, \frac{4}{m}\frac{2}{3}\frac{2}{m}$	$O_h$	48	Hexaoctahedron	Cube truncated by octahedron and by rhomb-dodecahedron	$m\bar{3}m$
Icosahedral system* (for details see Table 10.1.4.3)					
235 $m\bar{3}5, \frac{2}{m}\bar{3}5$	$I$ $I_h$	60 120	Pentagon-hexecontahedron Hecatonicosahedron	Snub pentagon-dodecahedron Pentagon-dodecahedron truncated by icosahedron and by rhomb-triacontahedron	– –
Cylindrical system†					
$\infty$ $\infty/m \equiv \overline{\infty}$ $\infty 2$	$C_\infty$ $C_{\infty h} \equiv S_\infty \equiv C_{\infty i}$ $D_\infty$	$\infty$ $\infty$ $\infty$	Rotating cone Rotating double cone 'Anti-rotating' double cone	Rotating circle Rotating finite cylinder 'Anti-rotating' finite cylinder	– – –
$\infty m$ $\infty/mm \equiv \overline{\infty}m, \frac{\infty 2}{m m} \equiv \overline{\infty}\frac{2}{m}$	$C_{\infty v}$ $D_{\infty h} \equiv D_{\infty d}$	$\infty$ $\infty$	Stationary cone Stationary double cone	Stationary circle Stationary finite cylinder	– –
Spherical system‡					
$2\infty$ $m\overline{\infty}, \frac{2}{m}\overline{\infty}$	$K$ $K_h$	$\infty$ $\infty$	Rotating sphere Stationary sphere	Rotating sphere Stationary sphere	– –

\* The Hermann–Mauguin symbols of the two icosahedral point groups are often written as 532 and  $\bar{5}3m$  (see text).

† Rotating and 'anti-rotating' forms in the cylindrical system have no 'vertical' mirror planes, whereas stationary forms have infinitely many vertical mirror planes. In classes  $\infty$  and  $\infty 2$ , enantiomorphism occurs, *i.e.* forms with opposite senses of rotation. Class  $\infty/m \equiv \overline{\infty}$  exhibits no enantiomorphism due to the centre of symmetry, even though the double cone is rotating in one direction. This can be understood as follows: One single rotating cone can be regarded as a right-handed or left-handed screw, depending on the sense of rotation with respect to the axial direction from the base to the tip of the cone. Thus, the rotating double cone consists of two cones with opposite handedness and opposite orientations related by the (single) horizontal mirror plane. In contrast, the 'anti-rotating' double cone in class  $\infty 2$  consists of two cones of equal handedness and opposite orientations, which are related by the (infinitely many) twofold axes. The term 'anti-rotating' means that upper and lower halves of the forms rotate in opposite directions.



‡ The spheres in class  $2\infty$  of the spherical system must rotate around an axis with at least two different orientations, in order to suppress all mirror planes. This class exhibits enantiomorphism, *i.e.* it contains spheres with either right-handed or left-handed senses of rotation around the axes (*cf.* Section 10.2.4, *Optical properties*). The stationary spheres in class  $m\overline{\infty}$  contain infinitely many mirror planes through the centres of the spheres.

Group  $2\infty$  is sometimes symbolized by  $\infty\infty$ ; group  $m\overline{\infty}$  by  $\overline{\infty}\overline{\infty}$  or  $\infty\infty m$ . The symbols used here indicate the minimal symmetry necessary to generate the groups; they show, furthermore, the relation to the cubic groups. The Schoenflies symbol  $K$  is derived from the German name *Kugelgruppe*.

10. POINT GROUPS AND CRYSTAL CLASSES

Table 10.1.4.3. *The two icosahedral point groups*

General, special and limiting face forms and point forms, oriented face- and site-symmetry symbols, and 'initial' values of  $(hkl)$  and  $x, y, z$  (see text).

235	$I$				
60	$d$	1	<p>Pentagon-hexecontahedron  <i>Snub pentagon-dodecahedron</i> (= <i>pentagon-dodecahedron</i> + <i>icosahedron</i> + <i>pentagon-hexecontahedron</i>)</p> <p>Trisicosahedron  <i>Pentagon-dodecahedron truncated by icosahedron</i>                      (poles between axes 2 and 3)</p> <p>Deltoid-hexecontahedron  <i>Rhomb-triacontahedron</i> &amp;  <i>pentagon-dodecahedron</i> &amp; <i>icosahedron</i>                      (poles between axes 3 and 5)</p> <p>Pentakisdodecahedron  <i>Icosahedron truncated by pentagon-dodecahedron</i>                      (poles between axes 5 and 2)</p>	<p><math>(hkl)</math>  <math>x, y, z</math></p> <p><math>(0kl)</math> with <math> l  &lt; 0.382 k </math>  <math>0, y, z</math> with <math> z  &lt; 0.382 y </math></p> <p><math>(0kl)</math> with <math>0.382 k  &lt;  l  &lt; 1.618 k </math>  <math>0, y, z</math> with <math>0.382 y  &lt;  z  &lt; 1.618 y </math></p> <p><math>(0kl)</math> with <math> l  &gt; 1.618 k </math>  <math>0, y, z</math> with <math> z  &gt; 1.618 y </math></p>	
30	$c$	2..	<p>Rhomb-triacontahedron  <i>Icosadodecahedron</i> (= <i>pentagon-dodecahedron</i> &amp; <i>icosahedron</i>)</p>	<p><math>(100)</math>  <math>x, 0, 0</math></p>	
20	$b$	.3.	<p>Regular icosahedron  <i>Regular pentagon-dodecahedron</i></p>	<p><math>(111)</math>  <math>x, x, x</math></p>	
12	$a$	..5	<p>Regular pentagon-dodecahedron  <i>Regular icosahedron</i></p>	<p><math>(01\tau)</math>  <math>0, y, \tau y</math> } with <math>\tau = \frac{1}{2}(\sqrt{5} + 1) = 1.618</math></p>	
Symmetry of special projections					
		Along $[001]$	Along $[111]$	Along $[1\tau 0]$	
		$2mm$	$3m$	$5m$	

10.1. CRYSTALLOGRAPHIC AND NONCRYSTALLOGRAPHIC POINT GROUPS

Table 10.1.4.3. *The two icosahedral point groups (cont.)*

$m\bar{3}5$	$I_h$			
				$\frac{2}{m}\bar{3}5$
120	<i>e</i>	1	Hecatonicosahedron or hexaicosahedron <i>Pentagon-dodecahedron truncated by icosahedron and by rhomb-triacontahedron</i>	$6 \blacksquare + 10 \blacktriangle + 15 \bullet + 15 m + \text{Centre}$ ( <i>hkl</i> ) <i>x, y, z</i>
60	<i>d</i>	<i>m..</i>	Trisicosahedron <i>Pentagon-dodecahedron truncated by icosahedron</i> (poles between axes 2 and $\bar{3}$ )	( <i>okl</i> ) with $ l  < 0.382 k $ <i>0, y, z</i> with $ z  < 0.382 y $
			Deltoid-hexecontahedron <i>Rhomb-triacontahedron &amp; pentagon-dodecahedron &amp; icosahedron</i> (poles between axes $\bar{3}$ and $\bar{5}$ )	( <i>okl</i> ) with $0.382 k  <  l  < 1.618 k $ <i>0, y, z</i> with $0.382 y  <  z  < 1.618 y $
			Pentakis-dodecahedron <i>Icosahedron truncated by pentagon-dodecahedron</i> (poles between axes $\bar{5}$ and 2)	( <i>okl</i> ) with $ l  > 1.618 k $ <i>0, y, z</i> with $ z  > 1.618 y $
30	<i>c</i>	$2mm..$	Rhomb-triacontahedron <i>Icosadodecahedron (= pentagon-dodecahedron &amp; icosahedron)</i>	(100) <i>x, 0, 0</i>
20	<i>b</i>	$3m (m\bar{3}.)$	Regular icosahedron <i>Regular pentagon-dodecahedron</i>	(111) <i>x, x, x</i>
12	<i>a</i>	$5m (m_5)$	Regular pentagon-dodecahedron <i>Regular icosahedron</i>	$(01\tau)$ $0, y, \tau y$ } with $\tau = \frac{1}{2}(\sqrt{5} + 1) = 1.618$
Symmetry of special projections				
		Along [001]	Along [111]	Along [1 $\tau$ 0]
		2 <i>mm</i>	6 <i>mm</i>	10 <i>mm</i>

10. POINT GROUPS AND CRYSTAL CLASSES

(b) the matrices  $Y^{-1}, Y^{-2}, Y^{-3}$  and  $Y^{-4}$  for the point coordinates.

This sequence of matrices ensures the same correspondence between the Miller indices and the point coordinates as for the crystallographic point groups in Table 10.1.2.2.

The matrices\* are

$$Y = Y^{-4} = \begin{pmatrix} \frac{1}{2} & g & G \\ g & G & -\frac{1}{2} \\ -G & \frac{1}{2} & g \end{pmatrix}, \quad Y^2 = Y^{-3} = \begin{pmatrix} -g & G & \frac{1}{2} \\ G & \frac{1}{2} & -g \\ -\frac{1}{2} & g & -G \end{pmatrix},$$

$$Y^3 = Y^{-2} = \begin{pmatrix} -g & G & -\frac{1}{2} \\ G & \frac{1}{2} & g \\ \frac{1}{2} & -g & -G \end{pmatrix}, \quad Y^4 = Y^{-1} = \begin{pmatrix} \frac{1}{2} & g & -G \\ g & G & \frac{1}{2} \\ G & -\frac{1}{2} & g \end{pmatrix},$$

with†

$$G = \frac{\sqrt{5} + 1}{4} = \frac{\tau}{2} = \cos 36^\circ = 0.80902 \simeq \frac{72}{89}$$

$$g = \frac{\sqrt{5} - 1}{4} = \frac{\tau - 1}{2} = \cos 72^\circ = 0.30902 \simeq \frac{17}{55}.$$

These matrices correspond to counter-clockwise rotations of 72, 144, 216 and 288° around a fivefold axis parallel to  $[1\tau 0]$ .

The resulting indices  $h, k, l$  and coordinates  $x, y, z$  are irrational but can be approximated closely by rational (or integral) numbers. This explains the occurrence of almost regular icosahedra or pentagon-dodecahedra as crystal forms (for instance pyrite) or atomic groups (for instance  $B_{12}$  icosahedron).


Further descriptions (including diagrams) of noncrystallographic groups are contained in papers by Nowacki (1933) and A. Niggli (1963) and in the textbooks by P. Niggli (1941, pp. 78–80, 96), Shubnikov & Koptsik (1974) and Vainshtein (1994). For the geometry of polyhedra, the well known books by H. S. M. Coxeter (especially Coxeter, 1973) are recommended.

10.1.4.3. Sub- and supergroups of the general point groups

In Figs. 10.1.4.1 to 10.1.4.3, the subgroup and supergroup relations between the two-dimensional and three-dimensional general point groups are illustrated. It should be remembered that the index of a group–subgroup relation between two groups of order infinity may be finite or infinite. For the two spherical groups, for instance, the index is  $[2]$ ; the cylindrical groups, on the other hand, are subgroups of index  $[\infty]$  of the spherical groups.

Fig. 10.1.4.1 for two dimensions shows that the two circular groups  $\infty m$  and  $\infty$  have subgroups of types  $nmm$  and  $n$ , respectively, each of index  $[\infty]$ . The order of the rotation point may be  $n = 4N, n = 4N + 2$  or  $n = 2N + 1$ . In the first case, the subgroups belong to the  $4N$ -gonal system, in the latter two cases, they belong to the  $(4N + 2)$ -gonal system. [In the diagram of the  $(4N + 2)$ -gonal system, the  $(2N + 1)$ -gonal groups appear with the symbols  $\frac{1}{2}nm$  and  $\frac{1}{2}n$ .] The subgroups of the circular groups are not maximal because for any given value of  $N$  there exists a group with  $N' = 2N$  which is both a subgroup of the circular group and a supergroup of the initial group.

The subgroup relations, for a specified value of  $N$ , within the  $4N$ -gonal and the  $(4N + 2)$ -gonal system, are shown in the lower part of the figure. They correspond to those of the crystallographic groups. A finite number of further maximal subgroups is obtained for lower



If  $n = 4N$ , go to  $4N$ -gonal system  
If  $n = 4N + 2$  or  $2N + 1$ , go to  $(4N + 2)$ -gonal system




Fig. 10.1.4.1. Subgroups and supergroups of the two-dimensional general point groups. Solid lines indicate maximal normal subgroups, double solid lines mean that there are two maximal normal subgroups with the same symbol. Dashed lines refer to sets of maximal conjugate subgroups. For the finite groups, the orders are given on the left. Note that the subgroups of the two circular groups are not maximal and the diagram applies only to a specified value of  $N$  (see text). For complete examples see Fig. 10.1.4.2.

values of  $N$ , until the crystallographic groups (Fig. 10.1.3.1) are reached. This is illustrated for the case  $N = 4$  in Fig. 10.1.4.2.

Fig. 10.1.4.3 for three dimensions illustrates that the two spherical groups  $2/m\infty$  and  $2\infty$  each have one infinite set of cylindrical maximal conjugate subgroups, as well as one infinite set




Fig. 10.1.4.2. The subgroups of the two-dimensional general point groups  $16mm$  ( $4N$ -gonal system) and  $18mm$  [ $(4N + 2)$ -gonal system including the  $(2N + 1)$ -gonal groups]. Compare with Fig. 10.1.4.1 which applies only to one value of  $N$ .

\* Note that for orthogonal matrices  $Y^{-1} = Y^t$  ( $t$  = transposed).  
† The number  $\tau = 2G = 2g + 1 = 1.618034$  (Fibonacci number) is the characteristic value of the golden section  $(\tau + 1) : \tau = \tau : 1$ , i.e.  $\tau(\tau - 1) = 1$ . Furthermore,  $\tau$  is the distance between alternating vertices of a regular pentagon of unit edge length and the distance from centre to vertex of a regular decagon of unit edge length.




Fig. 10.1.4.3. Subgroups and supergroups of the three-dimensional general point groups. Solid lines indicate maximal normal subgroups, double solid lines mean that there are two maximal normal subgroups with the same symbol. Dashed lines refer to sets of maximal conjugate subgroups. For the finite groups, the orders are given on the left and on the right. Note that the subgroups of the five cylindrical groups are not maximal and that the diagram applies only to a specified value of  $N$  (see text). Only those crystallographic point groups are included that are maximal subgroups of noncrystallographic point groups. Full Hermann–Mauguin symbols are used.

of cubic and one infinite set of icosahedral maximal finite conjugate subgroups, all of index  $[\infty]$ .

Each of the two icosahedral groups  $235$  and  $2/m\bar{3}5$  has one set of five cubic, one set of six pentagonal and one set of ten trigonal maximal conjugate subgroups of indices  $[5]$ ,  $[6]$  and  $[10]$ , respectively (*cf.* Section 10.1.4.2, *The two icosahedral groups*); they are listed on the right of Fig. 10.1.4.3. For crystallographic groups, Fig. 10.1.3.2 applies. The subgroup types of the five cylindrical point groups are shown on the left of Fig. 10.1.4.3. As explained above for two dimensions, these subgroups are *not maximal* and of index  $[\infty]$ . Depending upon whether the main symmetry axis has the multiplicity  $4N$ ,  $4N + 2$  or  $2N + 1$ , the subgroups belong to the  $4N$ -gonal,  $(4N + 2)$ -gonal or  $(2N + 1)$ -gonal system.

The subgroup and supergroup relations within these three systems are displayed in the lower left part of Fig 10.1.4.3. They are analogous to the crystallographic groups. To facilitate the use of

the diagrams, the  $(4N + 2)$ -gonal and the  $(2N + 1)$ -gonal systems are combined, with the consequence that the five classes of the  $(2N + 1)$ -gonal system now appear with the symbols  $\frac{1}{2}n\frac{2}{m}$ ,  $\frac{1}{2}n2$ ,  $\frac{1}{2}nm$ ,  $\frac{1}{2}n$  and  $\frac{1}{2}n$ . Again, the diagrams apply to a specified value of  $N$ . A finite number of further maximal subgroups is obtained for lower values of  $N$ , until the crystallographic groups (Fig. 10.1.3.2) are reached (*cf.* the two-dimensional examples in Fig. 10.1.4.2).

#### Acknowledgements

The authors are indebted to A. Niggli (Zürich) for valuable suggestions on Section 10.1.4, in particular for providing a sketch of Fig. 10.1.4.3. We thank H. Wondratschek (Karlsruhe) for stimulating and constructive discussions. We are grateful to R. A. Becker (Aachen) for the careful preparation of the diagrams.