

$I4_1/acd$

D_{4h}^{20}

$4/mmm$



Tetragonal

No. 142

$I 4_1/a 2/c 2/d$

Patterson symmetry $I4/mmm$

ORIGIN CHOICE 2



Origin at $\bar{1}$ at $b(c,a)d$, at $0, -\frac{1}{4}, \frac{1}{8}$ from $\bar{4}$

Asymmetric unit $0 \leq x \leq \frac{1}{2}; -\frac{1}{4} \leq y \leq \frac{1}{4}; 0 \leq z \leq \frac{1}{8}$

Symmetry operations

For $(0,0,0)+$ set

- | | | | |
|-----------------------------|--|---|---|
| (1) 1 | (2) $2(0,0,\frac{1}{2})$ $\frac{1}{4}, 0, z$ | (3) $4^+(0,0,\frac{1}{4})$ $-\frac{1}{4}, \frac{1}{2}, z$ | (4) $4^-(0,0,\frac{3}{4})$ $\frac{1}{4}, 0, z$ |
| (5) 2 $\frac{1}{4}, y, 0$ | (6) 2 $x, 0, \frac{1}{4}$ | (7) $2(\frac{1}{2}, \frac{1}{2}, 0)$ $x, x + \frac{1}{4}, \frac{3}{8}$ | (8) 2 $x, \bar{x} + \frac{1}{4}, \frac{1}{8}$ |
| (9) $\bar{1}$ $0, 0, 0$ | (10) a $x, y, \frac{1}{4}$ | (11) 4^+ $\frac{1}{2}, -\frac{1}{4}, z; \frac{1}{2}, -\frac{1}{4}, \frac{3}{8}$ | (12) 4^- $0, \frac{3}{4}, z; 0, \frac{3}{4}, \frac{1}{8}$ |
| (13) a $x, 0, z$ | (14) c $0, y, z$ | (15) $d(\frac{1}{4}, -\frac{1}{4}, \frac{1}{4})$ $x + \frac{1}{2}, \bar{x}, z$ | (16) $d(\frac{3}{4}, \frac{3}{4}, \frac{3}{4})$ x, x, z |

For $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})+$ set

- | | | | |
|---|--|---|---|
| (1) $i(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ | (2) 2 $0, \frac{1}{4}, z$ | (3) $4^+(0,0,\frac{3}{4})$ $\frac{1}{4}, \frac{1}{2}, z$ | (4) $4^-(0,0,\frac{1}{4})$ $\frac{3}{4}, 0, z$ |
| (5) $2(0, \frac{1}{2}, 0)$ $0, y, \frac{1}{4}$ | (6) $2(\frac{1}{2}, 0, 0)$ $x, \frac{1}{4}, 0$ | (7) $2(\frac{1}{2}, \frac{1}{2}, 0)$ $x, x - \frac{1}{4}, \frac{1}{8}$ | (8) 2 $x, \bar{x} + \frac{3}{4}, \frac{3}{8}$ |
| (9) $\bar{1}$ $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | (10) b $x, y, 0$ | (11) 4^+ $\frac{1}{2}, \frac{1}{4}, z; \frac{1}{2}, \frac{1}{4}, \frac{1}{8}$ | (12) 4^- $0, \frac{1}{4}, z; 0, \frac{1}{4}, \frac{3}{8}$ |
| (13) c $x, \frac{1}{4}, z$ | (14) b $\frac{1}{4}, y, z$ | (15) $d(-\frac{1}{4}, \frac{1}{4}, \frac{3}{4})$ $x + \frac{1}{2}, \bar{x}, z$ | (16) $d(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$ x, x, z |

Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; $t(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$; (2); (3); (5); (9)

Positions

Multiplicity, Wyckoff letter, Site symmetry	Coordinates		Reflection conditions			
	$(0,0,0) + (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}) +$		General:			
32 <i>g</i> 1	(1) x, y, z (5) $\bar{x} + \frac{1}{2}, y, \bar{z}$ (9) $\bar{x}, \bar{y}, \bar{z}$ (13) $x + \frac{1}{2}, \bar{y}, z$	(2) $\bar{x} + \frac{1}{2}, \bar{y}, z + \frac{1}{2}$ (6) $x, \bar{y}, \bar{z} + \frac{1}{2}$ (10) $x + \frac{1}{2}, y, \bar{z} + \frac{1}{2}$ (14) $\bar{x}, y, z + \frac{1}{2}$	(3) $\bar{y} + \frac{1}{4}, x + \frac{3}{4}, z + \frac{1}{4}$ (7) $y + \frac{1}{4}, x + \frac{3}{4}, \bar{z} + \frac{3}{4}$ (11) $y + \frac{3}{4}, \bar{x} + \frac{1}{4}, \bar{z} + \frac{3}{4}$ (15) $\bar{y} + \frac{3}{4}, \bar{x} + \frac{1}{4}, z + \frac{1}{4}$	(4) $y + \frac{1}{4}, \bar{x} + \frac{1}{4}, z + \frac{3}{4}$ (8) $\bar{y} + \frac{1}{4}, \bar{x} + \frac{1}{4}, \bar{z} + \frac{1}{4}$ (12) $\bar{y} + \frac{3}{4}, x + \frac{3}{4}, \bar{z} + \frac{1}{4}$ (16) $y + \frac{3}{4}, x + \frac{3}{4}, z + \frac{3}{4}$	$hkl : h + k + l = 2n$ $hk0 : h, k = 2n$ $0kl : k, l = 2n$ $hhl : 2h + l = 4n$ $00l : l = 4n$ $h00 : h = 2n$ $h\bar{h}0 : h = 2n$	
16 <i>f</i> ..2	$x, x + \frac{1}{4}, \frac{1}{8}$ $\bar{x}, \bar{x} + \frac{3}{4}, \frac{7}{8}$	$\bar{x} + \frac{1}{2}, \bar{x} + \frac{3}{4}, \frac{5}{8}$ $x + \frac{1}{2}, x + \frac{1}{4}, \frac{3}{8}$	$\bar{x}, x + \frac{3}{4}, \frac{3}{8}$ $x, \bar{x} + \frac{1}{4}, \frac{5}{8}$	$x + \frac{1}{2}, \bar{x} + \frac{1}{4}, \frac{7}{8}$ $\bar{x} + \frac{1}{2}, x + \frac{3}{4}, \frac{1}{8}$	$hkl : l = 2n + 1$ or $2h + l = 4n$	
16 <i>e</i> .2.	$x, 0, \frac{1}{4}$ $\bar{x}, 0, \frac{3}{4}$	$\bar{x} + \frac{1}{2}, 0, \frac{3}{4}$ $x + \frac{1}{2}, 0, \frac{1}{4}$	$\frac{1}{4}, x + \frac{3}{4}, \frac{1}{2}$ $\frac{3}{4}, \bar{x} + \frac{1}{4}, \frac{1}{2}$	$\frac{1}{4}, \bar{x} + \frac{1}{4}, 0$ $\frac{3}{4}, x + \frac{3}{4}, 0$	$hkl : l = 2n + 1$ or $h = 2n$	
16 <i>d</i> 2..	$0, \frac{1}{4}, z$ $0, \frac{3}{4}, \bar{z}$	$0, \frac{3}{4}, z + \frac{1}{4}$ $0, \frac{1}{4}, \bar{z} + \frac{3}{4}$	$\frac{1}{2}, \frac{1}{4}, \bar{z}$ $\frac{1}{2}, \frac{3}{4}, z$	$\frac{1}{2}, \frac{3}{4}, \bar{z} + \frac{3}{4}$ $\frac{1}{2}, \frac{1}{4}, z + \frac{1}{4}$	$hkl : 2h + l = 4n$	
16 <i>c</i> $\bar{1}$	0,0,0	$\frac{1}{2}, 0, \frac{1}{2}$	$\frac{1}{4}, \frac{3}{4}, \frac{1}{4}$ $\frac{1}{4}, \frac{1}{4}, \frac{3}{4}$	$\frac{1}{2}, 0, 0$ 0,0, $\frac{1}{2}$	$\frac{1}{4}, \frac{3}{4}, \frac{3}{4}$ $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}$	$hkl : h, k = 2n, h + k + l = 4n$
8 <i>b</i> 2.22	$0, \frac{1}{4}, \frac{1}{8}$	$0, \frac{3}{4}, \frac{3}{8}$	$0, \frac{3}{4}, \frac{7}{8}$	$0, \frac{1}{4}, \frac{5}{8}$	$hkl : 2h + l = 4n$	
8 <i>a</i> $\bar{4}$..	$0, \frac{1}{4}, \frac{3}{8}$	$0, \frac{3}{4}, \frac{5}{8}$	$\frac{1}{2}, \frac{1}{4}, \frac{5}{8}$	$\frac{1}{2}, \frac{3}{4}, \frac{3}{8}$	$hkl : 2h + l = 4n$	

Symmetry of special projections

Along [001] $p4mm$
 $\mathbf{a}' = \frac{1}{2}\mathbf{a}$ $\mathbf{b}' = \frac{1}{2}\mathbf{b}$
Origin at $\frac{1}{4}, 0, z$

Along [100] $p2mm$
 $\mathbf{a}' = \frac{1}{2}\mathbf{b}$ $\mathbf{b}' = \frac{1}{2}\mathbf{c}$
Origin at $x, 0, 0$

Along [110] $c2mm$
 $\mathbf{a}' = \frac{1}{2}(-\mathbf{a} + \mathbf{b})$ $\mathbf{b}' = \frac{1}{2}\mathbf{c}$
Origin at $x, x + \frac{1}{4}, \frac{1}{8}$

Maximal non-isomorphic subgroups

I	[2] $I\bar{4}2d$ (122)	(1; 2; 5; 6; 11; 12; 15; 16)+
	[2] $I\bar{4}c2$ (120)	(1; 2; 7; 8; 11; 12; 13; 14)+
	[2] $I4, cd$ (110)	(1; 2; 3; 4; 13; 14; 15; 16)+
	[2] $I4, 22$ (98)	(1; 2; 3; 4; 5; 6; 7; 8)+
	[2] $I4_1/a11$ ($I4_1/a, 88$)	(1; 2; 3; 4; 9; 10; 11; 12)+
	[2] $I2/a2/c1$ ($Ibca, 73$)	(1; 2; 5; 6; 9; 10; 13; 14)+
	[2] $I2/a12/d$ ($Fddd, 70$)	(1; 2; 7; 8; 9; 10; 15; 16)+

IIa none**IIb** none**Maximal isomorphic subgroups of lowest index****IIc** [3] $I4_1/acd$ ($\mathbf{c}' = 3\mathbf{c}$) (142); [9] $I4_1/acd$ ($\mathbf{a}' = 3\mathbf{a}, \mathbf{b}' = 3\mathbf{b}$) (142)**Minimal non-isomorphic supergroups****I** [3] $Fd\bar{3}c$ (228); [3] $Ia\bar{3}d$ (230)**II** [2] $C4_2/amd$ ($\mathbf{c}' = \frac{1}{2}\mathbf{c}$) ($P4_2/nm, 134$)