

$R\bar{3}m$

D_{3d}^5

$\bar{3}m$



Trigonal

No. 166

$R\bar{3}2/m$

Patterson symmetry $R\bar{3}m$

HEXAGONAL AXES



Origin at centre ($\bar{3}m$)

Asymmetric unit $0 \leq x \leq \frac{2}{3}$; $0 \leq y \leq \frac{2}{3}$; $0 \leq z \leq \frac{1}{6}$; $x \leq 2y$; $y \leq \min(1-x, 2x)$

Vertices $0, 0, 0$ $\frac{2}{3}, \frac{1}{3}, 0$ $\frac{1}{3}, \frac{2}{3}, 0$
 $0, 0, \frac{1}{6}$ $\frac{2}{3}, \frac{1}{3}, \frac{1}{6}$ $\frac{1}{3}, \frac{2}{3}, \frac{1}{6}$

Symmetry operations

For (0,0,0)+ set

- | | | |
|----------------------|------------------------------|------------------------------|
| (1) 1 | (2) $3^+ 0,0,z$ | (3) $3^- 0,0,z$ |
| (4) $2 x,x,0$ | (5) $2 x,0,0$ | (6) $2 0,y,0$ |
| (7) $\bar{1} 0,0,0$ | (8) $\bar{3}^+ 0,0,z; 0,0,0$ | (9) $\bar{3}^- 0,0,z; 0,0,0$ |
| (10) $m x,\bar{x},z$ | (11) $m x,2x,z$ | (12) $m 2x,x,z$ |

For $(\frac{2}{3}, \frac{1}{3}, \frac{1}{3})$ + set

- | | | |
|--|--|--|
| (1) $t(\frac{2}{3}, \frac{1}{3}, \frac{1}{3})$ | (2) $3^+(0,0,\frac{1}{3}) \frac{1}{3}, \frac{1}{3}, z$ | (3) $3^-(0,0,\frac{1}{3}) \frac{1}{3}, 0, z$ |
| (4) $2(\frac{1}{2}, \frac{1}{2}, 0) x, x - \frac{1}{6}, \frac{1}{6}$ | (5) $2(\frac{1}{2}, 0, 0) x, \frac{1}{6}, \frac{1}{6}$ | (6) $2 \frac{1}{3}, y, \frac{1}{6}$ |
| (7) $\bar{1} \frac{1}{3}, \frac{1}{6}, \frac{1}{6}$ | (8) $\bar{3}^+ \frac{1}{3}, -\frac{1}{3}, z; \frac{1}{3}, -\frac{1}{3}, \frac{1}{6}$ | (9) $\bar{3}^- \frac{1}{3}, \frac{2}{3}, z; \frac{1}{3}, \frac{2}{3}, \frac{1}{6}$ |
| (10) $g(\frac{1}{6}, -\frac{1}{6}, \frac{1}{3}) x + \frac{1}{2}, \bar{x}, z$ | (11) $g(\frac{1}{6}, \frac{1}{3}, \frac{1}{3}) x + \frac{1}{4}, 2x, z$ | (12) $g(\frac{2}{3}, \frac{1}{3}, \frac{1}{3}) 2x, x, z$ |

For $(\frac{1}{3}, \frac{2}{3}, \frac{2}{3})$ + set

- | | | |
|--|--|--|
| (1) $t(\frac{1}{3}, \frac{2}{3}, \frac{2}{3})$ | (2) $3^+(0,0,\frac{2}{3}) 0, \frac{1}{3}, z$ | (3) $3^-(0,0,\frac{2}{3}) \frac{1}{3}, \frac{1}{3}, z$ |
| (4) $2(\frac{1}{2}, \frac{1}{2}, 0) x, x + \frac{1}{6}, \frac{1}{3}$ | (5) $2 x, \frac{1}{3}, \frac{1}{3}$ | (6) $2(0, \frac{1}{2}, 0) \frac{1}{6}, y, \frac{1}{3}$ |
| (7) $\bar{1} \frac{1}{6}, \frac{1}{3}, \frac{1}{3}$ | (8) $\bar{3}^+ \frac{2}{3}, \frac{1}{3}, z; \frac{2}{3}, \frac{1}{3}, \frac{1}{3}$ | (9) $\bar{3}^- -\frac{1}{3}, \frac{1}{3}, z; -\frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ |
| (10) $g(-\frac{1}{6}, \frac{1}{6}, \frac{2}{3}) x + \frac{1}{2}, \bar{x}, z$ | (11) $g(\frac{1}{3}, \frac{2}{3}, \frac{2}{3}) x, 2x, z$ | (12) $g(\frac{1}{3}, \frac{1}{6}, \frac{2}{3}) 2x - \frac{1}{2}, x, z$ |

Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; $t(\frac{2}{3}, \frac{1}{3}, \frac{1}{3})$; (2); (4); (7)

Positions

Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

(0,0,0)+ $(\frac{2}{3}, \frac{1}{3}, \frac{1}{3})$ + $(\frac{1}{3}, \frac{2}{3}, \frac{2}{3})$ +

Reflection conditions

General:

- $hkil : -h + k + l = 3n$
 $hki0 : -h + k = 3n$
 $hh\bar{2}hl : l = 3n$
 $h\bar{h}0l : h + l = 3n$
 $000l : l = 3n$
 $h\bar{h}00 : h = 3n$

Special: no extra conditions

36	<i>i</i>	1	(1) x, y, z	(2) $\bar{y}, x - y, z$	(3) $\bar{x} + y, \bar{x}, z$	(4) y, x, \bar{z}	(5) $x - y, \bar{y}, \bar{z}$	(6) $\bar{x}, \bar{x} + y, \bar{z}$	(7) $\bar{x}, \bar{y}, \bar{z}$	(8) $y, \bar{x} + y, \bar{z}$	(9) $x - y, x, \bar{z}$	(10) \bar{y}, \bar{x}, z	(11) $\bar{x} + y, y, z$	(12) $x, x - y, z$
18	<i>h</i>	. <i>m</i>	x, \bar{x}, z	$x, 2x, z$	$2\bar{x}, \bar{x}, z$	\bar{x}, x, \bar{z}	$2x, x, \bar{z}$	$\bar{x}, 2\bar{x}, \bar{z}$						
18	<i>g</i>	. 2	$x, 0, \frac{1}{2}$	$0, x, \frac{1}{2}$	$\bar{x}, \bar{x}, \frac{1}{2}$	$\bar{x}, 0, \frac{1}{2}$	$0, \bar{x}, \frac{1}{2}$	$x, x, \frac{1}{2}$						
18	<i>f</i>	. 2	$x, 0, 0$	$0, x, 0$	$\bar{x}, \bar{x}, 0$	$\bar{x}, 0, 0$	$0, \bar{x}, 0$	$x, x, 0$						
9	<i>e</i>	. $2/m$	$\frac{1}{2}, 0, 0$	$0, \frac{1}{2}, 0$	$\frac{1}{2}, \frac{1}{2}, 0$									
9	<i>d</i>	. $2/m$	$\frac{1}{2}, 0, \frac{1}{2}$	$0, \frac{1}{2}, \frac{1}{2}$	$\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$									
6	<i>c</i>	3 <i>m</i>	$0, 0, z$	$0, 0, \bar{z}$										
3	<i>b</i>	$\bar{3} m$	$0, 0, \frac{1}{2}$											
3	<i>a</i>	$\bar{3} m$	$0, 0, 0$											

Symmetry of special projections

Along [001] $p6mm$

$\mathbf{a}' = \frac{1}{3}(2\mathbf{a} + \mathbf{b})$ $\mathbf{b}' = \frac{1}{3}(-\mathbf{a} + \mathbf{b})$
 Origin at 0, 0, z

Along [100] $p2$

$\mathbf{a}' = \frac{1}{2}(\mathbf{a} + 2\mathbf{b})$ $\mathbf{b}' = \frac{1}{3}(-\mathbf{a} - 2\mathbf{b} + \mathbf{c})$
 Origin at x, 0, 0

Along [210] $p2mm$

$\mathbf{a}' = \frac{1}{2}\mathbf{b}$ $\mathbf{b}' = \frac{1}{3}\mathbf{c}$
 Origin at $x, \frac{1}{2}x, 0$

HEXAGONAL AXES

Maximal non-isomorphic subgroups

- I** [2] $R\bar{3}m$ (160) (1; 2; 3; 10; 11; 12)+
 [2] $R\bar{3}2$ (155) (1; 2; 3; 4; 5; 6)+
 [2] $R\bar{3}1$ ($R\bar{3}$, 148) (1; 2; 3; 7; 8; 9)+
 { [3] $R12/m$ ($C2/m$, 12) (1; 4; 7; 10)+
 [3] $R12/m$ ($C2/m$, 12) (1; 5; 7; 11)+
 [3] $R12/m$ ($C2/m$, 12) (1; 6; 7; 12)+
- IIa** { [3] $P\bar{3}m1$ (164) 1; 2; 3; 4; 5; 6; 7; 8; 9; 10; 11; 12
 [3] $P\bar{3}m1$ (164) 1; 2; 3; 10; 11; 12; (4; 5; 6; 7; 8; 9) + $(\frac{2}{3}, \frac{1}{3}, \frac{1}{3})$
 [3] $P\bar{3}m1$ (164) 1; 2; 3; 10; 11; 12; (4; 5; 6; 7; 8; 9) + $(\frac{1}{3}, \frac{2}{3}, \frac{2}{3})$
- IIb** [2] $R\bar{3}c$ ($\mathbf{a}' = -\mathbf{a}, \mathbf{b}' = -\mathbf{b}, \mathbf{c}' = 2\mathbf{c}$) (167)

Maximal isomorphic subgroups of lowest index

- IIc** [2] $R\bar{3}m$ ($\mathbf{a}' = -\mathbf{a}, \mathbf{b}' = -\mathbf{b}, \mathbf{c}' = 2\mathbf{c}$) (166); [4] $R\bar{3}m$ ($\mathbf{a}' = -2\mathbf{a}, \mathbf{b}' = -2\mathbf{b}$) (166)

Minimal non-isomorphic supergroups

- I** [4] $Pm\bar{3}m$ (221); [4] $Pn\bar{3}m$ (224); [4] $Fm\bar{3}m$ (225); [4] $Fd\bar{3}m$ (227); [4] $Im\bar{3}m$ (229)
- II** [3] $P\bar{3}1m$ ($\mathbf{a}' = \frac{1}{3}(2\mathbf{a} + \mathbf{b}), \mathbf{b}' = \frac{1}{3}(-\mathbf{a} + \mathbf{b}), \mathbf{c}' = \frac{1}{3}\mathbf{c}$) (162)

RHOMBOHEDRAL AXES

Maximal non-isomorphic subgroups

- I** [2] $R\bar{3}m$ (160) 1; 2; 3; 10; 11; 12
 [2] $R\bar{3}2$ (155) 1; 2; 3; 4; 5; 6
 [2] $R\bar{3}1$ ($R\bar{3}$, 148) 1; 2; 3; 7; 8; 9
 { [3] $R12/m$ ($C2/m$, 12) 1; 4; 7; 10
 [3] $R12/m$ ($C2/m$, 12) 1; 5; 7; 11
 [3] $R12/m$ ($C2/m$, 12) 1; 6; 7; 12
- IIa** none
- IIb** [2] $F\bar{3}c$ ($\mathbf{a}' = 2\mathbf{a}, \mathbf{b}' = 2\mathbf{b}, \mathbf{c}' = 2\mathbf{c}$) ($R\bar{3}c$, 167); [3] $P\bar{3}m1$ ($\mathbf{a}' = \mathbf{a} - \mathbf{b}, \mathbf{b}' = \mathbf{b} - \mathbf{c}, \mathbf{c}' = \mathbf{a} + \mathbf{b} + \mathbf{c}$) (164)

Maximal isomorphic subgroups of lowest index

- IIc** [2] $R\bar{3}m$ ($\mathbf{a}' = \mathbf{b} + \mathbf{c}, \mathbf{b}' = \mathbf{a} + \mathbf{c}, \mathbf{c}' = \mathbf{a} + \mathbf{b}$) (166); [4] $R\bar{3}m$ ($\mathbf{a}' = -\mathbf{a} + \mathbf{b} + \mathbf{c}, \mathbf{b}' = \mathbf{a} - \mathbf{b} + \mathbf{c}, \mathbf{c}' = \mathbf{a} + \mathbf{b} - \mathbf{c}$) (166)

Minimal non-isomorphic supergroups

- I** [4] $Pm\bar{3}m$ (221); [4] $Pn\bar{3}m$ (224); [4] $Fm\bar{3}m$ (225); [4] $Fd\bar{3}m$ (227); [4] $Im\bar{3}m$ (229)
- II** [3] $P\bar{3}1m$ ($\mathbf{a}' = \frac{1}{3}(2\mathbf{a} - \mathbf{b} - \mathbf{c}), \mathbf{b}' = \frac{1}{3}(-\mathbf{a} + 2\mathbf{b} - \mathbf{c}), \mathbf{c}' = \frac{1}{3}(\mathbf{a} + \mathbf{b} + \mathbf{c})$) (162)

Trigonal

$\bar{3}m$

D_{3d}^5

$R\bar{3}m$

Patterson symmetry $R\bar{3}m$

$R\bar{3}2/m$

No. 166

RHOMBOHEDRAL AXES
(For drawings see hexagonal axes)

Origin at centre ($\bar{3}m$)

Asymmetric unit $0 \leq x \leq 1; 0 \leq y \leq 1; 0 \leq z \leq \frac{1}{2}; y \leq x; z \leq \min(y, 1-x)$
Vertices $0,0,0 \quad 1,0,0 \quad 1,1,0 \quad \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$

Symmetry operations

- | | | |
|-----------------------|----------------------------------|----------------------------------|
| (1) 1 | (2) $3^+ x, x, x$ | (3) $3^- x, x, x$ |
| (4) $2 \bar{x}, 0, x$ | (5) $2 x, \bar{x}, 0$ | (6) $2 0, y, \bar{y}$ |
| (7) $\bar{1} 0, 0, 0$ | (8) $\bar{3}^+ x, x, x; 0, 0, 0$ | (9) $\bar{3}^- x, x, x; 0, 0, 0$ |
| (10) $m x, y, x$ | (11) $m x, x, z$ | (12) $m x, y, y$ |

Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (4); (7)

Positions

Multiplicity, Wyckoff letter, Site symmetry	Coordinates	Reflection conditions
12 <i>i</i> 1	(1) x, y, z (2) z, x, y (3) y, z, x (4) $\bar{z}, \bar{y}, \bar{x}$ (5) $\bar{y}, \bar{x}, \bar{z}$ (6) $\bar{x}, \bar{z}, \bar{y}$ (7) $\bar{x}, \bar{y}, \bar{z}$ (8) $\bar{z}, \bar{x}, \bar{y}$ (9) $\bar{y}, \bar{z}, \bar{x}$ (10) z, y, x (11) y, x, z (12) x, z, y	General: no conditions
6 <i>h</i> . <i>m</i>	x, x, z z, x, x x, z, x $\bar{z}, \bar{x}, \bar{x}$ $\bar{x}, \bar{x}, \bar{z}$ $\bar{x}, \bar{z}, \bar{x}$	Special: no extra conditions
6 <i>g</i> .2	$x, \bar{x}, \frac{1}{2}$ $\frac{1}{2}, x, \bar{x}$ $\bar{x}, \frac{1}{2}, x$ $\bar{x}, x, \frac{1}{2}$ $\frac{1}{2}, \bar{x}, x$ $x, \frac{1}{2}, \bar{x}$	
6 <i>f</i> .2	$x, \bar{x}, 0$ $0, x, \bar{x}$ $\bar{x}, 0, x$ $\bar{x}, x, 0$ $0, \bar{x}, x$ $x, 0, \bar{x}$	
3 <i>e</i> . $2/m$	$0, \frac{1}{2}, \frac{1}{2}$ $\frac{1}{2}, 0, \frac{1}{2}$ $\frac{1}{2}, \frac{1}{2}, 0$	
3 <i>d</i> . $2/m$	$\frac{1}{2}, 0, 0$ $0, \frac{1}{2}, 0$ $0, 0, \frac{1}{2}$	
2 <i>c</i> 3 <i>m</i>	x, x, x $\bar{x}, \bar{x}, \bar{x}$	
1 <i>b</i> $\bar{3}m$	$\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$	
1 <i>a</i> $\bar{3}m$	$0, 0, 0$	

Symmetry of special projections

Along $[111] p6mm$ $\mathbf{a}' = \frac{1}{3}(2\mathbf{a} - \mathbf{b} - \mathbf{c})$ $\mathbf{b}' = \frac{1}{3}(-\mathbf{a} + 2\mathbf{b} - \mathbf{c})$ Origin at x, x, x	Along $[1\bar{1}0] p2$ $\mathbf{a}' = \frac{1}{2}(\mathbf{a} + \mathbf{b} - 2\mathbf{c})$ $\mathbf{b}' = \mathbf{c}$ Origin at $x, \bar{x}, 0$	Along $[2\bar{1}\bar{1}] p2mm$ $\mathbf{a}' = \frac{1}{2}(\mathbf{b} - \mathbf{c})$ $\mathbf{b}' = \frac{1}{3}(\mathbf{a} + \mathbf{b} + \mathbf{c})$ Origin at $2x, \bar{x}, \bar{x}$
---	--	--

(Continued on preceding page)