

Tetragonal

6. SCANNING TABLES

 Laue class $C_{4h} - 4/m$
Laue class $C_{4h} - 4/m$
Geometric class $C_4 - 4$

 No. 75 $P4$

$$\mathcal{G} = P4$$

 C_4^1

Orientation orbit (<i>hkl</i>)	Conventional basis of the scanning group a' b' d	Scanning group \mathcal{H}	Linear orbit sd	Sectional layer group $\mathcal{L}(sd)$	
(001)	a b c	$P4$	sd	$p4$	L49

 No. 76 $P4_1$

$$\mathcal{G} = P4_1$$

 C_4^2

Orientation orbit (<i>hkl</i>)	Conventional basis of the scanning group a' b' d	Scanning group \mathcal{H}	Linear orbit sd	Sectional layer group $\mathcal{L}(sd)$	
(001)	a b c	$P4_1$	$[sd, (s + \frac{1}{4})d, (s + \frac{1}{2})d, (s + \frac{3}{4})d]$	$p1$	L01

 No. 77 $P4_2$

$$\mathcal{G} = P4_2$$

 C_4^3

Orientation orbit (<i>hkl</i>)	Conventional basis of the scanning group a' b' d	Scanning group \mathcal{H}	Linear orbit sd	Sectional layer group $\mathcal{L}(sd)$	
(001)	a b c	$P4_2$	$[sd, (s + \frac{1}{2})d]$	$p112$	L03

 No. 78 $P4_3$

$$\mathcal{G} = P4_3$$

 C_4^4

Orientation orbit (<i>hkl</i>)	Conventional basis of the scanning group a' b' d	Scanning group \mathcal{H}	Linear orbit sd	Sectional layer group $\mathcal{L}(sd)$	
(001)	a b c	$P4_3$	$[sd, (s + \frac{1}{4})d, (s + \frac{1}{2})d, (s + \frac{3}{4})d]$	$p1$	L01

No. 88 $I4_1/a$

$\mathcal{G} = I4_1/a$ origin 2

C_{4h}^6

Orientation orbit (<i>hkl</i>)	Conventional basis of the scanning group			Scanning group \mathcal{H}	Linear orbit <i>sd</i>	Sectional layer group	
	<i>a'</i>	<i>b'</i>	<i>d</i>			$\mathcal{L}(sd)$	
(001)	a	b	c	$I4_1/a$ (origin 2)	$[0\mathbf{d}, \frac{1}{2}\mathbf{d};$ $\frac{1}{4}\mathbf{d}, \frac{3}{4}\mathbf{d}]$ $[\frac{1}{8}\mathbf{d}, \frac{5}{8}\mathbf{d};$ $\frac{3}{8}\mathbf{d}, \frac{7}{8}\mathbf{d}]$ $[\pm s\mathbf{d}, (\pm s + \frac{1}{4})\mathbf{d},$ $(\pm s + \frac{1}{2})\mathbf{d}, (\pm s + \frac{3}{4})\mathbf{d}]$	$p112/b$	L07
						$p112/a [(\mathbf{a} + \mathbf{b})/4]$	L07
						$p\bar{4} (\mathbf{b}/4)$	L50
						$p\bar{4} (3\mathbf{b}/4)$	L50
						$p112 (\mathbf{b}/4)$	L03

Auxiliary tables for Laue class $C_{4h} - 4/m$

Centring types *P* and *I*

Orientation orbit (<i>hkl</i>)	Conventional basis of the scanning group			Auxiliary basis of the scanning group		
	<i>a'</i>	<i>b'</i>	<i>d</i>	$\hat{\mathbf{a}}$	$\hat{\mathbf{b}}$	$\hat{\mathbf{c}}$
(<i>mn</i> 0)	c	$n\mathbf{a} - m\mathbf{b}$	$p\mathbf{a} + q\mathbf{b}$	a	b	c
(\bar{m} n0)	c	$m\mathbf{a} + n\mathbf{b}$	$-q\mathbf{a} + p\mathbf{b}$			

Arithmetic classes $4P$ and $4I$

Serial No.	75	76	77	78	79	80
Group type	C_4^1	C_4^2	C_4^3	C_4^4	C_4^5	C_4^6
Group	$P4$	$P4_1$	$P4_2$	$P4_3$	$I4$	$I4_1$
(<i>mn</i> 0)	$P112$	$P112_1$	$P112$	$P112_1$	$I112$	$I112$
(\bar{m} n0)						

Arithmetic classes $\bar{4}P$ and $\bar{4}$

Serial No.	81	82
Group type	S_4^1	S_4^2
Group	$P\bar{4}$	$I\bar{4}$
(<i>mn</i> 0)	$P112$	$I112$
(\bar{m} n0)		

Arithmetic class $4/mP$

Serial No.	83	84	85		86	
			Origin 1	Origin 2	Origin 1	Origin 2
Group type	C_{4h}^1	C_{4h}^2	C_{4h}^3		C_{4h}^4	
Group	$P4/m$	$P4_2/m$	$P4/n$		$P4_2/n$	
(<i>mn</i> 0)	$P112/m$	$P112/m$	$P112/n$	$P112/n$	$P112/n$	$P112/n$
(\bar{m} n0)			$(\mathbf{a} + \mathbf{b})/4$		$(\mathbf{a} + \mathbf{b} + \mathbf{c})/4$	