

## REFERENCES

## 22.1 (cont.)

- Richards, F. M. (1985). Calculation of molecular volumes and areas for structures of known geometry. *Methods Enzymol.* **115**, 440–464.
- Richards, F. M. & Lim, W. A. (1994). An analysis of packing in the protein folding problem. *Q. Rev. Biophys.* **26**, 423–498.
- Richmond, T. J. (1984). Solvent accessible surface area and excluded volume in proteins: analytical equations for overlapping spheres and implications for the hydrophobic effect. *J. Mol. Biol.* **178**, 63–89.
- Richmond, T. J. & Richards, F. M. (1978). Packing of alpha-helices: geometrical constraints and contact areas. *J. Mol. Biol.* **119**, 537–555.
- Rossmann, M. G. (1989). The canyon hypothesis. *J. Biol. Chem.* **264**, 14587–14590.
- Rossmann, M. G. & Palmenberg, A. C. (1988). Conservation of the putative receptor attachment site in picornaviruses. *Virology*, **164**, 373–382.
- Rowland, R. S. & Taylor, R. (1996). Intermolecular nonbonded contact distances in organic crystal structures: comparison with distances expected from van der Waals radii. *J. Phys. Chem.* **100**, 7384–7391.
- Sgro, J.-Y. (1996). Virus visualization. In *Encyclopedia of virology plus* (CD-ROM version), edited by R. G. Webster & A. Granoff. San Diego: Academic Press.
- Sharp, K. A., Nicholls, A., Fine, R. F. & Honig, B. (1991). Reconciling the magnitude of the microscopic and macroscopic hydrophobic effects. *Science*, **252**, 107–109.
- Sherry, B., Mosser, A. G., Colonna, R. J. & Rueckert, R. R. (1986). Use of monoclonal antibodies to identify four neutralization immunogens on a common cold picornavirus, human rhinovirus 14. *J. Virol.* **57**, 246–257.
- Sherry, B. & Rueckert, R. (1985). Evidence for at least two dominant neutralization antigens on human rhinovirus 14. *J. Virol.* **53**, 137–143.
- Shrake, A. & Rupley, J. A. (1973). Environment and exposure to solvent of protein atoms. *Lysozyme and insulin*. *J. Mol. Biol.* **79**, 351–371.
- Sibbald, P. R. & Argos, P. (1990). Weighting aligned protein or nucleic acid sequences to correct for unequal representation. *J. Mol. Biol.* **216**, 813–818.
- Singh, R. K., Tropsha, A. & Vaisman, I. I. (1996). Delaunay tessellation of proteins: four body nearest-neighbor propensities of amino acid residues. *J. Comput. Biol.* **3**, 213–222.
- Sreenivasan, U. & Axelsen, P. H. (1992). Buried water in homologous serine proteases. *Biochemistry*, **31**, 12785–12791.
- Tanford, C. (1997). How protein chemists learned about the hydrophobicity factor. *Protein Sci.* **6**, 1358–1366.
- Tanford, C. H. (1979). Interfacial free energy and the hydrophobic effect. *Proc. Natl Acad. Sci. USA*, **76**, 4175–4176.
- Ten Eyck, L. F. (1977). Efficient structure-factor calculation for large molecules by the fast Fourier transform. *Acta Cryst.* **A33**, 486–492.
- Tsai, J., Gerstein, M. & Levitt, M. (1996). Keeping the shape but changing the charges: a simulation study of urea and its isosteric analogues. *J. Chem. Phys.* **104**, 9417–9430.
- Tsai, J., Gerstein, M. & Levitt, M. (1997). Estimating the size of the minimal hydrophobic core. *Protein Sci.* **6**, 2606–2616.
- Tsai, J., Taylor, R., Chothia, C. & Gerstein, M. (1999). The packing density in proteins: standard radii and volumes. *J. Mol. Biol.* **290**, 253–266.
- Tsai, J., Voss, N. & Gerstein, M. (2001). Voronoi calculations of protein volumes: sensitivity analysis and parameter database. *Bioinformatics*. In the press.
- Voronoi, G. F. (1908). Nouvelles applications des paramètres continus à la théorie des formes quadratiques. *J. Reine Angew. Math.* **134**, 198–287.
- Williams, M. A., Goodfellow, J. M. & Thornton, J. M. (1994). Buried waters and internal cavities in monomeric proteins. *Protein Sci.* **3**, 1224–1235.
- Wodak, S. J. & Janin, J. (1980). Analytical approximation to the accessible surface areas of proteins. *Proc. Natl Acad. Sci. USA*, **77**, 1736–1740.
- Xie, Q. & Chapman, M. S. (1996). Canine parvovirus capsid structure, analyzed at 2.9 Å resolution. *J. Mol. Biol.* **264**, 497–520.
- Zhou, G., Somasundaram, T., Blanc, E., Parthasarathy, G., Ellington, W. R. & Chapman, M. S. (1998). Transition state structure of arginine kinase: implications for catalysis of bimolecular reactions. *Proc. Natl Acad. Sci. USA*, **95**, 8449–8454.

## 22.2

- Adman, E., Watenpaugh, K. D. & Jensen, L. H. (1975). N—H...S hydrogen bonds in *Peptococcus aerogenes ferredoxin*, *Clostridium pasteurianum rubredoxin* and *Chromatium high potential iron protein*. *Proc. Natl Acad. Sci. USA*, **72**, 4854–4858.
- Alber, T., Dao-pin, S., Wilson, K., Wozniak, J. A., Cook, S. P. & Matthews, B. W. (1987). Contributions of hydrogen bonds of Thr 157 to the thermodynamic stability of phage T4 lysozyme. *Nature (London)*, **330**, 41–46.
- Artymiuk, P. J. & Blake, C. C. F. (1981). Refinement of human lysozyme at 1.5 Å resolution. Analysis of non-bonded and hydrogen-bonded interactions. *J. Mol. Biol.* **152**, 737–762.
- Baker, E. N. (1995). Solvent interactions with proteins as revealed by X-ray crystallographic studies. In *Protein-solvent interactions*, edited by R. B. Gregory, pp. 143–189. New York: Marcel Dekker Inc.
- Baker, E. N. & Hubbard, R. E. (1984). Hydrogen bonding in globular proteins. *Prog. Biophys. Mol. Biol.* **44**, 97–179.
- Blundell, T., Barlow, D., Borkakoti, N. & Thornton, J. (1983). Solvent-induced distortions and the curvature of  $\alpha$ -helices. *Nature (London)*, **306**, 281–283.
- Bordo, D. & Argos, P. (1994). The role of side-chain hydrogen bonds in the formation and stabilization of secondary structure in soluble proteins. *J. Mol. Biol.* **243**, 504–519.
- Burley, S. K. & Petsko, G. A. (1986). Amino-aromatic interactions in proteins. *FEBS Lett.* **203**, 139–143.
- Cate, J. H., Gooding, A. R., Podell, E., Zhou, K., Golden, B. L., Kundrot, C. E., Cech, T. R. & Doudna, J. A. (1996). Crystal structure of a group I ribozyme domain: principles of RNA packing. *Science*, **273**, 1678–1685.
- Derewenda, Z. S., Derewenda, U. & Kobos, P. (1994). (His)C $\epsilon$ —H...O=C< hydrogen bond in the active site of serine hydrolases. *J. Mol. Biol.* **241**, 83–93.
- Derewenda, Z. S., Lee, L. & Derewenda, U. (1995). The occurrence of C—H...O hydrogen bonds in proteins. *J. Mol. Biol.* **252**, 248–262.
- Edwards, R. A., Baker, H. M., Whittaker, M. M., Whittaker, J. W. & Baker, E. N. (1998). Crystal structure of *Escherichia coli* manganese superoxide dismutase at 2.1 Å resolution. *J. Biol. Inorg. Chem.* **3**, 161–171.
- Fersht, A. R. & Serrano, L. (1993). Principles in protein stability derived from protein engineering experiments. *Curr. Opin. Struct. Biol.* **3**, 75–83.
- Fersht, A. R., Shi, J.-P., Knill-Jones, J., Lowe, D. M., Wilkinson, A. J., Blow, D. M., Brick, P., Carter, P., Waye, M. M. Y. & Winter, G. (1985). Hydrogen bonding and biological specificity analysed by protein engineering. *Nature (London)*, **314**, 235–238.
- Flocco, M. M. & Mowbray, S. L. (1995). Strange bedfellows: interactions between acidic side-chains in proteins. *J. Mol. Biol.* **254**, 96–105.
- Gregoret, L. M., Rader, S. D., Fletterick, R. J. & Cohen, F. E. (1991). Hydrogen bonds involving sulfur atoms in proteins. *Proteins Struct. Funct. Genet.* **9**, 99–107.
- Hagler, A. T., Huler, E. & Lifson, S. (1974). Energy functions for peptides and proteins. I. Derivation of a consistent force field including the hydrogen bond from amide crystals. *J. Am. Chem. Soc.* **96**, 5319–5327.
- Harper, E. T. & Rose, G. D. (1993). Helix stop signals in proteins and peptides: the capping box. *Biochemistry*, **32**, 7605–7609.
- Huggins, M. L. (1971). 50 years of hydrogen bonding theory. *Angew. Chem. Int. Ed. Engl.* **10**, 147–208.

## 22.2 (cont.)

- Ippolito, J. A., Alexander, R. S. & Christianson, D. W. (1990). *Hydrogen bond stereochemistry in protein structure and function*. *J. Mol. Biol.* **215**, 457–471.
- Jeffrey, G. A. & Saenger, W. (1991). *Hydrogen bonding in biological structures*. New York: Springer-Verlag.
- Kauzmann, W. (1959). *Some factors in the interpretation of protein denaturation*. *Adv. Protein Chem.* **14**, 1–64.
- Ligon, A. C. & Millen, D. J. (1987). *Directional character, strength, and nature of the hydrogen bond in gas-phase dimers*. *Acc. Chem. Res.* **20**, 39–45.
- Levitt, M. & Perutz, M. F. (1988). *Aromatic rings act as hydrogen bond acceptors*. *J. Mol. Biol.* **201**, 751–754.
- McDonald, I. K. & Thornton, J. M. (1994a). *Satisfying hydrogen bonding potential in proteins*. *J. Mol. Biol.* **238**, 777–793.
- McDonald, I. K. & Thornton, J. M. (1994b). *The application of hydrogen bonding analysis in X-ray crystallography to help orientate asparagine, glutamine and histidine side chains*. *Protein Eng.* **8**, 217–224.
- Matthews, B. W. (1972). *The  $\gamma$  turn. Evidence for a new folded conformation in proteins*. *Macromolecules*, **5**, 818–819.
- Mitchell, J. B. O., Nandi, C. L., McDonald, I. K., Thornton, J. M. & Price, S. L. (1994). *Amino/aromatic interactions in proteins: is the evidence stacked against hydrogen bonding?* *J. Mol. Biol.* **239**, 315–331.
- Nemethy, G. & Printz, M. P. (1972). *The  $\gamma$  turn, a possible folded conformation of the polypeptide chain. Comparison with the  $\beta$  turn*. *Macromolecules*, **5**, 755–758.
- Pauling, L. (1960). *The nature of the chemical bond*, 3rd ed. Ithaca: Cornell University Press.
- Pauling, L. & Corey, R. B. (1951). *Configurations of polypeptide chains with favoured orientations around single bonds: two new pleated sheets*. *Proc. Natl Acad. Sci. USA*, **37**, 729–740.
- Pauling, L., Corey, R. B. & Branson, H. R. (1951). *The structure of proteins: two hydrogen-bonded helical configurations of the polypeptide chain*. *Proc. Natl Acad. Sci. USA*, **37**, 205–211.
- Pley, H. W., Flaherty, K. M. & McKay, D. B. (1994). *Three-dimensional structure of a hammerhead ribozyme*. *Nature (London)*, **372**, 68–74.
- Presta, L. G. & Rose, G. D. (1988). *Helix signals in proteins*. *Science*, **240**, 1632–1641.
- Richardson, J. S., Getzoff, E. D. & Richardson, D. C. (1978). *The  $\beta$ -bulge: a common small unit of nonrepetitive protein structure*. *Proc. Natl Acad. Sci. USA*, **75**, 2574–2578.
- Richardson, J. S. & Richardson, D. C. (1988). *Amino acid preferences for specific locations at the ends of  $\alpha$ -helices*. *Science*, **240**, 1648–1652.
- Savage, H. J., Elliott, C. J., Freeman, C. M. & Finney, J. L. (1993). *Lost hydrogen bonds and buried surface area: rationalising stability in globular proteins*. *J. Chem. Soc. Faraday Trans.* **89**, 2609–2617.
- Schellman, C. (1980). *The alpha-L conformation at the ends of helices*. In *Protein folding*, edited by R. Jaenicke, pp. 53–61. Amsterdam: Elsevier.
- Stickle, D. F., Presta, L. G., Dill, K. A. & Rose, G. D. (1992). *Hydrogen bonding in globular proteins*. *J. Mol. Biol.* **226**, 1143–1159.
- Sutor, D. J. (1962). *The C—H...O hydrogen bond in crystals*. *Nature (London)*, **195**, 68–69.
- Taylor, R., Kennard, O. & Versichel, W. (1983). *Geometry of the N—H...O=C hydrogen bond. I. Lone pair directionality*. *J. Am. Chem. Soc.* **105**, 5761–5766.
- Thanki, N., Thornton, J. M. & Goodfellow, J. M. (1988). *Distribution of water around amino acids in proteins*. *J. Mol. Biol.* **202**, 637–657.
- Thanki, N., Umrana, Y., Thornton, J. M. & Goodfellow, J. M. (1991). *Analysis of protein main-chain solvation as a function of secondary structure*. *J. Mol. Biol.* **221**, 669–691.
- Wahl, M. C. & Sundaralingam, M. (1997). *C—H...O hydrogen bonding in biology*. *Trends Biochem. Sci.* **22**, 97–102.
- Williams, M. A., Goodfellow, J. M. & Thornton, J. M. (1994). *Buried waters and internal cavities in monomeric proteins*. *Protein Sci.* **3**, 1224–1235.

## 22.3

- Antosiewicz, J., McCammon, J. A. & Gilson, M. K. (1994). *Prediction of pH-dependent properties of proteins*. *J. Mol. Biol.* **238**, 415–436.
- Åqvist, J., Luecke, H., Quijoco, F. A. & Warshel, A. (1991). *Dipoles localized at helix termini of proteins stabilize charges*. *Proc. Natl Acad. Sci. USA*, **88**, 2026–2030.
- Bacquet, R. & Rossky, P. (1984). *Ionic atmosphere of rodlike polyelectrolytes. A hypernetted chain study*. *J. Phys. Chem.* **88**, 2660.
- Bashford, D. & Karplus, M. (1990). *pK<sub>a</sub>'s of ionizable groups in proteins: atomic detail from a continuum electrostatic model*. *Biochemistry*, **29**, 10219–10225.
- Beroza, P., Fredkin, D., Okamura, M. & Feher, G. (1991). *Protonation of interacting residues in a protein by a Monte-Carlo method*. *Proc. Natl Acad. Sci. USA*, **88**, 5804–5808.
- Bharadwaj, R., Windemuth, A., Sridharan, S., Honig, B. & Nicholls, A. (1994). *The fast multipole boundary-element method for molecular electrostatics – an optimal approach for large systems*. *J. Comput. Chem.* **16**, 898–913.
- Bruccoleri, R. E., Novotny, J., Sharp, K. A. & Davis, M. E. (1996). *Finite difference Poisson–Boltzmann electrostatic calculations: increased accuracy achieved by harmonic dielectric smoothing and charge anti-aliasing*. *J. Comput. Chem.* **18**, 268–276.
- Davis, M. E. (1994). *The inducible multipole solvation model – a new model for solvation effects on solute electrostatics*. *J. Chem. Phys.* **100**, 5149–5159.
- Davis, M. E. & McCammon, J. A. (1989). *Solving the finite difference linear Poisson Boltzmann equation: comparison of relaxational and conjugate gradient methods*. *J. Comput. Chem.* **10**, 386–395.
- Gilson, M. (1993). *Multiple-site titration and molecular modeling: 2. Rapid methods for computing energies and forces for ionizable groups in proteins*. *Proteins Struct. Funct. Genet.* **15**, 266–282.
- Gilson, M., Davis, M., Luty, B. & McCammon, J. (1993). *Computation of electrostatic forces on solvated molecules using the Poisson–Boltzmann equation*. *J. Phys. Chem.* **97**, 3591–3600.
- Gilson, M. & Honig, B. (1986). *The dielectric constant of a folded protein*. *Biopolymers*, **25**, 2097–2119.
- Gilson, M., McCammon, J. & Madura, J. (1995). *Molecular dynamics simulation with continuum solvent*. *J. Comput. Chem.* **16**, 1081–1095.
- Gilson, M., Sharp, K. A. & Honig, B. (1988). *Calculating the electrostatic potential of molecules in solution: method and error assessment*. *J. Comput. Chem.* **9**, 327–335.
- Holst, M., Kozack, R., Saied, F. & Subramaniam, S. (1994). *Protein electrostatics – rapid multigrid-based Newton algorithm for solution of the full nonlinear Poisson–Boltzmann equation*. *J. Biomol. Struct. Dyn.* **11**, 1437–1445.
- Holst, M. & Saied, F. (1993). *Multigrid solution of the Poisson–Boltzmann equation*. *J. Comput. Chem.* **14**, 105–113.
- Jayaram, B., Fine, R., Sharp, K. A. & Honig, B. (1989). *Free energy calculations of ion hydration: an analysis of the Born model in terms of microscopic simulations*. *J. Phys. Chem.* **93**, 4320–4327.
- Jayaram, B., Sharp, K. A. & Honig, B. (1989). *The electrostatic potential of B-DNA*. *Biopolymers*, **28**, 975–993.
- Jean-Charles, A., Nicholls, A., Sharp, K., Honig, B., Tempczyk, A., Hendrickson, T. & Still, C. (1990). *Electrostatic contributions to solvation energies: comparison of free energy perturbation and continuum calculations*. *J. Am. Chem. Soc.* **113**, 1454–1455.
- Langsetmo, K., Fuchs, J. A., Woodward, C. & Sharp, K. A. (1991). *Linkage of thioredoxin stability to titration of ionizable groups with perturbed pK<sub>a</sub>*. *Biochemistry*, **30**, 7609–7614.
- Lee, F., Chu, Z. & Warshel, A. (1993). *Microscopic and semimicroscopic calculations of electrostatic energies in proteins by the Polaris and Enzymix programs*. *J. Comput. Chem.* **14**, 161–185.