

$D_4^7$  $P4_322$ 

No. 95

 $P4_322$ **Generators selected** (1);  $t(1,0,0)$ ;  $t(0,1,0)$ ;  $t(0,0,1)$ ; (2); (3); (5)**General position**

Multiplicity,  
Wyckoff letter,  
Site symmetry

8	$d$	1	(1) $x,y,z$ (2) $\bar{x},\bar{y},z + \frac{1}{2}$ (3) $\bar{y},x,z + \frac{3}{4}$ (4) $y,\bar{x},z + \frac{1}{4}$ (5) $\bar{x},y,\bar{z}$ (6) $x,\bar{y},\bar{z} + \frac{1}{2}$ (7) $y,x,\bar{z} + \frac{1}{4}$ (8) $\bar{y},\bar{x},\bar{z} + \frac{3}{4}$	Coordinates
---	-----	---	--	-------------

**I Maximal translationengleiche subgroups**

[2] $P4_311$ (78, $P4_3$ )	1; 2; 3; 4		
[2] $P2_112$ (20, $C222_1$ )	1; 2; 7; 8	$\mathbf{a} - \mathbf{b}, \mathbf{a} + \mathbf{b}, \mathbf{c}$	0, 0, 3/8
[2] $P2_121$ (17, $P222_1$ )	1; 2; 5; 6		0, 0, 1/4

**II Maximal klassengleiche subgroups****• Enlarged unit cell**

[2] $\mathbf{a}' = 2\mathbf{a}$ , $\mathbf{b}' = 2\mathbf{b}$			
$C4_322_1$ (96, $P4_32_12$ )	$\langle 2; 5; 3 + (1, 0, 0) \rangle$	$\mathbf{a} - \mathbf{b}, \mathbf{a} + \mathbf{b}, \mathbf{c}$	0, 0, 1/4
$C4_322_1$ (96, $P4_32_12$ )	$\langle 2; 3; 5 + (1, 0, 0) \rangle$	$\mathbf{a} - \mathbf{b}, \mathbf{a} + \mathbf{b}, \mathbf{c}$	1/2, 1/2, 1/4
$C4_322$ (95, $P4_322$ )	$\langle 2; 3; 5 + (0, 0, 1) \rangle$	$\mathbf{a} - \mathbf{b}, \mathbf{a} + \mathbf{b}, \mathbf{c}$	0, 0, 1/8
$C4_322$ (95, $P4_322$ )	$\langle 2 + (1, 1, 0); 3 + (1, 0, 0); 5 + (1, 0, 1) \rangle$	$\mathbf{a} - \mathbf{b}, \mathbf{a} + \mathbf{b}, \mathbf{c}$	1/2, 1/2, 1/8
[3] $\mathbf{c}' = 3\mathbf{c}$			
$\begin{cases} P4_122 \text{ (91)} \\ P4_122 \text{ (91)} \\ P4_122 \text{ (91)} \end{cases}$	$\begin{cases} \langle 3; 5; 2 + (0, 0, 1) \rangle \\ \langle 3; 2 + (0, 0, 1); 5 + (0, 0, 2) \rangle \\ \langle 3; 2 + (0, 0, 1); 5 + (0, 0, 4) \rangle \end{cases}$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	0, 0, 1
		$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	0, 0, 2

**• Series of maximal isomorphic subgroups**

[p] $\mathbf{c}' = p\mathbf{c}$			
$P4_322$ (95)	$\langle 2 + (0, 0, \frac{p}{2} - \frac{1}{2}); 3 + (0, 0, \frac{3p}{4} - \frac{3}{4}); 5 + (0, 0, 2u) \rangle$ $p > 4; 0 \leq u < p$	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	0, 0, $u$
$P4_122$ (91)	$\langle 2 + (0, 0, \frac{p}{2} - \frac{1}{2}); 3 + (0, 0, \frac{p}{4} - \frac{3}{4}); 5 + (0, 0, 2u) \rangle$ $p > 2; 0 \leq u < p$	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	0, 0, $u$
$[p^2]$ $\mathbf{a}' = p\mathbf{a}$ , $\mathbf{b}' = p\mathbf{b}$			
$P4_322$ (95)	$\langle 2 + (2u, 2v, 0); 3 + (u + v, -u + v, 0); 5 + (2u, 0, 0) \rangle$ $p > 2; 0 \leq u < p; 0 \leq v < p$	$p\mathbf{a}, p\mathbf{b}, \mathbf{c}$	$u, v, 0$
	$p^2$ conjugate subgroups for the prime $p$		

**I Minimal translationengleiche supergroups**

none

**II Minimal non-isomorphic klassengleiche supergroups****• Additional centring translations**[2]  $I4_122$  (98)**• Decreased unit cell**[2]  $\mathbf{c}' = \frac{1}{2}\mathbf{c}$   $P4_222$  (93)