

4.2. DISORDER DIFFUSE SCATTERING OF X-RAYS AND NEUTRONS

$$F_l(\mathbf{H}) = \sum_j \sum_n f_j J_n(2\pi H_r r_j) \exp\{-in\psi_j\} \\ \times \exp(2\pi i l z_j) \exp\{in[\Psi + (\pi/2)]\} \quad (4.2.4.35)$$

or

$$F_l(\mathbf{H}) = \sum_n (\alpha_n + i\beta_n) \exp\{in\Psi\} \\ \alpha_n = \sum_j f_j J_n(2\pi H_r r_j) \cos\{n[(\pi/2) - \psi_j] + 2\pi l z_j\} \\ \beta_n = \sum_j f_j J_n(2\pi H_r r_j) \sin\{n[(\pi/2) - \psi_j] + 2\pi l z_j\}.$$

Intensity in the l th diffuse layer is given by

$$I_l = \sum_n \sum_{n'} [(\alpha_n \alpha_{n'} + \beta_n \beta_{n'}) + i(\alpha_{n'} \beta_n - \alpha_n \beta_{n'})] \\ \times \exp\{i(n - n')\Psi\}. \quad (4.2.4.36)$$

(a) *Cylinder symmetry* (free rotating molecules around the chain axis or statistical averaging with respect to ψ over an assembly of chains). Only component F_{0l} occurs:

$$F_{0l}(H_r, L) = 2\pi \int \int \langle a_M \rangle J_0(2\pi H_r r) \exp\{2\pi i l z\} r \, dr \, dz$$

or

$$F_{0l}(H_r, L) = \sum_j f_j J_0(2\pi H_r r_j) \exp\{2\pi i l z_j\}.$$

In particular, $F_{00}(H_r)$ determines the radial component of the molecule projected along z :

$$F_{00}(H_r) = \sum_j f_j J_0(2\pi H_r r_j).$$

(b) *p-fold symmetry* of a plane molecule (or projected molecule) as outlined previously: only components np instead of n occur. Bessel functions J_0 and J_p are sufficient in most cases.

(c) *Vertical mirror plane*: see above.

(d) *Horizontal mirror plane* (perpendicular to the chain): Exponentials $\exp\{2\pi i l z\}$ in equation (4.2.4.32) may be replaced by $\cos 2\pi l z$.

(e) *Twofold symmetry axis* perpendicular to the chain axis (at positions $\psi = 0, 2\pi/p, \dots$). Exponentials in equation (4.2.4.32), $\exp\{-i(np\psi - 2\pi l z)\}$, are replaced by the corresponding cosine term $\cos(np\psi + 2\pi l z)$.

Formulae concerning the reverse method (Fourier synthesis) are not given here (see, e.g., Vainshtein, 1966). Usually there is no practical use in diffuse-scattering work because it is very difficult to separate out a single component F_{nl} . Every diffuse layer is affected by *all* components F_{nl} . There is a chance if *one* diffuse layer corresponds predominantly to *one* Bessel function.

4.2.4.3.2. Disorder within randomly distributed collinear chains

Deviations from strict periodicities in the z direction within one chain may be due to loss of translational symmetry of the centres of the molecules along z and/or due to varying orientations of the molecules with respect to different axes, such as azimuthal misorientation, tilting with respect to the z axis or combinations of both types. As in 3D crystals, there may or may not exist 1D structures in an averaged sense.

4.2.4.3.2.1. General treatment

All formulae given in this section are only special cases of a 3D treatment (see, e.g., Guinier, 1963). The 1D lattice (4.2.4.12) is

replaced by a distribution:

$$d(z) = \sum_\nu \delta(z - z_\nu) \\ D(L) = \sum_\nu \exp\{2\pi i L z_\nu\} \quad (4.2.4.37)$$

$$F(\mathbf{H}) = F_M(\mathbf{H})D(L).$$

The Patterson function is given by

$$P(\mathbf{r}) = [a_M(\mathbf{r}) * a_M - (\mathbf{r})] * [d(z) * d - (z)]. \quad (4.2.4.38)$$

Because the autocorrelation function $w = d * d$ is centrosymmetric

$$w(z) = N\delta(z) + \sum_\nu \sum_\mu \delta[z - (z_\nu - z_\mu)] + \sum_\nu \sum_\mu \delta[z + (z_\nu - z_\mu)], \quad (4.2.4.39)$$

the interference function $W(L) (= |D(L)|^2)$ is given by

$$W(L) = N + 2 \sum_\nu \sum_\mu \cos 2\pi [L(z_\nu - z_\mu)] \quad (4.2.4.40)$$

$$I(\mathbf{H}) = |F_M(\mathbf{H})|^2 W(L). \quad (4.2.4.41)$$

Sometimes, e.g. in the following example of orientational disorder, there is an order only within domains. As shown in Section 4.2.3, this may be treated by a box or shape function $b(z) = 1$ for $z \leq z_N$ and 0 elsewhere.

$$d(z) = d_\infty b(z) \\ a(\mathbf{r}) = a_M(\mathbf{r}) * [d_\infty b(z)] \quad (4.2.4.42)$$

$$F(\mathbf{H}) = F_M(\mathbf{H}) [D_\infty * B(L)]$$

with

$$b(z) * b - (z) \leftrightarrow |B(L)|^2 \\ I = |F_M(\mathbf{H})|^2 |D_\infty * B(L)|^2. \quad (4.2.4.43)$$

If the order is perfect within one domain one has $D_\infty(L) \simeq \sum \delta(L - l)$; $(D_\infty * B) = \sum D(L - l)$; i.e. each reflection is affected by the shape function.

4.2.4.3.2.2. Orientational disorder

A misorientation of the chain molecules with respect to one another is taken into account by different structure factors F_M .

$$I(\mathbf{H}) = \sum_\nu \sum_\mu F_\nu(\mathbf{H}) F_\mu(\mathbf{H})^+ \exp\{2\pi i L(z_\nu - z_\mu)\}. \quad (4.2.4.44)$$

A further discussion follows the same arguments outlined in Section 4.2.3. For example, a very simple result is found in the case of uncorrelated orientations. Averaging over all pairs $F_\nu F_\mu^+$ yields

$$I(\mathbf{H}) = N(\langle |F|^2 \rangle - \langle |F| \rangle^2) + \langle |F| \rangle^2 L(L), \quad (4.2.4.44a)$$

where

$$\langle |F| \rangle^2 = 1/N^2 \langle F_\nu F_\mu^+ \rangle \\ = \sum_\nu \alpha_\nu F_\nu(\mathbf{H}) \sum_\mu \alpha_\mu F_\mu^+(\mathbf{H}) \quad (\nu \neq \mu) \\ \langle |F|^2 \rangle = 1/N \langle F_\nu F_\nu^+ \rangle = \sum_\nu \alpha_\nu |F_\nu(\mathbf{H})|^2.$$

Besides the diffuse layer system there is a diffuse background modulated by the \mathbf{H} dependence of $[\langle |F(\mathbf{H})|^2 \rangle - \langle |F(\mathbf{H})| \rangle^2]$.