

1.1. INTRODUCTION TO THE PROPERTIES OF TENSORS

Furthermore, the left-hand term of (1.1.4.11) remains unchanged if we interchange the indices  $i$  and  $j$ . The terms on the right-hand side therefore also remain unchanged, whatever the value of  $T_{ll}$  or  $T_{kl}$ . It follows that

$$s_{ijll} = s_{jill}$$

$$s_{ijkl} = s_{ijlk} = s_{jikl} = s_{jilk}$$

Similar relations hold for  $c_{ijkl}$ ,  $Q_{ijkl}$ ,  $p_{ijkl}$  and  $\pi_{ijkl}$ : the submatrices **2** and **3**, **4** and **7**, **5**, **6**, **8** and **9**, respectively, are equal.

Equation (1.4.1.11) can be rewritten, introducing the coefficients of the Voigt strain matrix:

$$S_\alpha = S_{ii} = \sum_l s_{iill} T_{ll} + \sum_{k \neq l} (s_{iikl} + s_{iilk}) T_{kl} \quad (\alpha = 1, 2, 3)$$

$$S_\alpha = S_{ij} + S_{ji} = \sum_l (s_{ijll} + s_{jill}) T_{ll} + \sum_{k \neq l} (s_{ijkl} + s_{ijlk} + s_{jikl} + s_{jilk}) T_{kl} \quad (\alpha = 4, 5, 6).$$

We shall now introduce a two-index notation for the elastic compliances, according to the following conventions:

$$\left. \begin{aligned} i = j; \quad k = l; \quad s_{\alpha\beta} &= s_{iill} \\ i = j; \quad k \neq l; \quad s_{\alpha\beta} &= s_{iikl} + s_{iilk} \\ i \neq j; \quad k = l; \quad s_{\alpha\beta} &= s_{ijkk} + s_{jikl} \\ i \neq j; \quad k \neq l; \quad s_{\alpha\beta} &= s_{ijkl} + s_{jikl} + s_{ijlk} + s_{jilk} \end{aligned} \right\} \quad (1.1.4.12)$$

We have thus associated with the fourth-rank tensor a square  $6 \times 6$  matrix with 36 coefficients:

$\beta$	1	2	3	4	5	6
$\alpha$						
1	11	12	13	14	15	16
2	21	22	23	24	25	26
3	31	32	33	34	35	36
4	41	42	43	44	45	46
5	51	52	53	54	55	56
6	61	62	63	64	65	66

One can translate relation (1.1.4.12) using the  $9 \times 9$  matrix representing  $s_{ijkl}$  by adding term by term the coefficients of submatrices **2** and **3**, **4** and **7** and **5**, **6**, **8** and **9**, respectively:

$$\left( \begin{array}{c} 1 \\ 2+3 \end{array} \right) = \left( \begin{array}{c|c} 1 & 2 \\ \hline 4+7 & 5+6 \\ & +8+9 \end{array} \right) \times \left( \begin{array}{c} 1 \\ 2+3 \end{array} \right)$$

Using the two-index notation, equation (1.1.4.9) becomes

$$S_\alpha = s_{\alpha\beta} T_\beta \quad (1.1.4.13)$$

A similar development can be applied to the other fourth-rank tensors  $\pi_{ijkl}$ , which will be replaced by  $6 \times 6$  matrices with 36 coefficients, according to the following rules.

(i) *Elastic stiffnesses*,  $c_{ijkl}$  and *elasto-optic coefficients*,  $p_{ijkl}$ :

$$\left( \begin{array}{c} 1 \\ 2 \end{array} \right) = \left( \begin{array}{c|c} 1 & 2 \\ \hline 4 & 5 \end{array} \right) \times \left( \begin{array}{c} 1 \\ 2 \end{array} \right)$$

where

$$c_{\alpha\beta} = c_{ijkl}$$

$$p_{\alpha\beta} = p_{ijkl}$$

(ii) *Piezo-optic coefficients*,  $\pi_{ijkl}$ :

$$\left( \begin{array}{c} 1 \\ 2 \end{array} \right) = \left( \begin{array}{c|c} 1 & 2+3 \\ \hline 4 & 5+6 \end{array} \right) \times \left( \begin{array}{c} 1 \\ 2 \end{array} \right)$$


where

$$\left. \begin{aligned} i = j; \quad k = l; \quad \pi_{\alpha\beta} &= \pi_{iill} \\ i = j; \quad k \neq l; \quad \pi_{\alpha\beta} &= \pi_{iikl} + \pi_{iilk} \\ i \neq j; \quad k = l; \quad \pi_{\alpha\beta} &= \pi_{ijkk} = \pi_{jikl} \\ i \neq j; \quad k \neq l; \quad \pi_{\alpha\beta} &= \pi_{ijkl} + \pi_{jikl} = \pi_{ijlk} + \pi_{jilk} \end{aligned} \right\}$$

(iii) *Electrostriction coefficients*,  $Q_{ijkl}$ : same relation as for the elastic compliances.


1.1.4.10.6. Independent components of the matrix associated with a fourth-rank tensor according to the following point groups

1.1.4.10.6.1. Triclinic system, groups  $\bar{1}$ , 1




1.1.4.10.6.2. Monoclinic system

Groups  $2/m$ ,  $2$ ,  $m$ , twofold axis parallel to  $Ox_2$ :



1.1.4.10.6.3. Orthorhombic system




Groups  $mmm$ ,  $2mm$ ,  $222$ :



# 1. TENSORIAL ASPECTS OF PHYSICAL PROPERTIES



## 1.1.4.10.6.4. Trigonal system

(i) Groups 3,  $\bar{3}$ :




## 1.1.4.10.6.5. Tetragonal system

(i) Groups 4,  $\bar{4}$  and 4/m:




(ii) Groups 422, 4mm,  $\bar{4}2m$  and 4/mmm:





where  $\ominus$  is a component numerically equal but opposite in sign to the heavy dot component to which it is linked;  $\oplus$  is a component equal to twice the heavy dot component to which it is linked;  $\odot$  is a component equal to minus twice the heavy dot component to which it is linked;  $\otimes$  is equal to  $1/2(p_{11} - p_{12})$ ,  $(\pi_{11} - \pi_{12})$ ,  $2(Q_{11} - Q_{12})$ ,  $1/2(c_{11} - c_{12})$  and  $2(s_{11} - s_{12})$ , respectively.

(ii) Groups 32, 3m,  $\bar{3}m$ :




## 1.1.4.10.6.6. Hexagonal system

(i) Groups 6,  $\bar{6}$  and 6/m:




1.1. INTRODUCTION TO THE PROPERTIES OF TENSORS




1.1.4.10.6.8. Spherical system

For all tensors



(ii) Groups 622, 6mm,  $\bar{6}2m$  and 6/mmm:



1.1.4.10.7. Reduction of the number of independent components of axial tensors of rank 2

It was shown in Section 1.1.4.5.3.2 that axial tensors of rank 2 are actually tensors of rank 3 antisymmetric with respect to two indices. The matrix of independent components of a tensor such that

$$g_{ijk} = -g_{jik}$$

is given by

$$\left( \begin{array}{ccc|ccc|cc} & 122 & 133 & 123 & 131 & & 132 & & 121 \\ -121 & & 223 & & 231 & -122 & 232 & -123 & \\ -131 & -232 & & -233 & & -132 & & -133 & -231 \end{array} \right).$$

The second-rank axial tensor  $g_{kl}$  associated with this tensor is defined by


$$g_{kl} = \frac{1}{2} \epsilon_{ijk} g_{ijl}$$

For instance, the piezomagnetic coefficients that give the magnetic moment  $M_i$  due to an applied stress  $T_\alpha$  are the components of a second-rank axial tensor,  $\Lambda_{i\alpha}$  (see Section 1.5.7.1):

$$M_i = \Lambda_{i\alpha} T_\alpha$$

1.1.4.10.6.7. Cubic system


(i) Groups 23 and 3m:



1.1.4.10.7.1. Independent components according to the following point groups

(i) Triclinic system


(a) Group 1:



(b) Group  $\bar{1}$ : all components are equal to zero.

(ii) Monoclinic system

(a) Group 2:



(ii) Groups 432,  $\bar{4}3m$  and  $m\bar{3}m$ :

