

$\bar{3}1c$

$31m$



Trigonal

No. 50

$\bar{3}1c$

Patterson symmetry $\bar{3}1m$

SECOND SETTING



Origin on $31c$

Asymmetric unit $0 \leq x; 0 \leq y; 0 \leq z \leq \frac{1}{2}$

Symmetry operations

- | | | |
|-----------------|-----------------|-----------------|
| (1) 1 | (2) $3^+ 0,0,z$ | (3) $3^- 0,0,z$ |
| (4) $c \ x,x,z$ | (5) $c \ x,0,z$ | (6) $c \ 0,y,z$ |

Generators selected (1); $t(0,0,1)$; (2); (4)

Positions

Multiplicity, Wyckoff letter, Site symmetry	Coordinates			Reflection conditions
				General:
6 <i>b</i> 1	(1) x, y, z (4) $y, x, z + \frac{1}{2}$	(2) $\bar{y}, x - y, z$ (5) $x - y, \bar{y}, z + \frac{1}{2}$	(3) $\bar{x} + y, \bar{x}, z$ (6) $\bar{x}, \bar{x} + y, z + \frac{1}{2}$	$l : l = 2n$
2 <i>a</i> 3..	0, 0, z	0, 0, $z + \frac{1}{2}$		Special: no extra conditions

Symmetry of special projections

Along [001] $3m$	Along [100] $\bar{3}11g$	Along [210] $\bar{3}1$
Origin at 0, 0, z	$\mathbf{a}' = \mathbf{c}$ Origin at $x, 0, 0$	$\mathbf{a}' = \frac{1}{2}\mathbf{c}$ Origin at $x, \frac{1}{2}x, 0$

Maximal non-isotypic non-enantiomorphic subgroups

I	$[2]\bar{3}11$ ($\bar{3}3, 42$)	1; 2; 3
	$[3]\bar{3}11c$ ($\bar{3}c11, 5$)	1; 4
	$[3]\bar{3}11c$ ($\bar{3}c11, 5$)	1; 5
	$[3]\bar{3}11c$ ($\bar{3}c11, 5$)	1; 6

IIa none

IIb none

Maximal isotypic subgroups and enantiomorphic subgroups of lowest index

IIc $[3]\bar{3}11c$ ($\mathbf{c}' = 3\mathbf{c}$) ($\bar{3}3c1, 50$)

Minimal non-isotypic non-enantiomorphic supergroups

I $[2]\bar{3}11c$ (52); $[2]\bar{3}6cc$ (69); $[2]\bar{3}6_3mc$ (70); $[2]\bar{3}6c2$ (72)

II $[2]\bar{3}m1$ ($\mathbf{c}' = \frac{1}{2}\mathbf{c}$) (49)