

Tetragonal

6. SCANNING TABLES

Laue class $C_{4h} - 4/m$ **Laue class $C_{4h} - 4/m$** **Geometric class $C_4 - 4$** No. 75 $P4$

$$\mathcal{G} = P4$$

 C_4^1

Orientation orbit (hkl)	Conventional basis of the scanning group $\mathbf{a}' \quad \mathbf{b}' \quad \mathbf{d}$	Scanning group \mathcal{H}	Linear orbit $s\mathbf{d}$	Sectional layer group $\mathcal{L}(s\mathbf{d})$
(001)	$\mathbf{a} \quad \mathbf{b} \quad \mathbf{c}$	$P4$	$s\mathbf{d}$	$p4$ L49

No. 76 $P4_1$

$$\mathcal{G} = P4_1$$

 C_4^2

Orientation orbit (hkl)	Conventional basis of the scanning group $\mathbf{a}' \quad \mathbf{b}' \quad \mathbf{d}$	Scanning group \mathcal{H}	Linear orbit $s\mathbf{d}, (s + \frac{1}{4})\mathbf{d},$ $(s + \frac{1}{2})\mathbf{d}, (s + \frac{3}{4})\mathbf{d}$	Sectional layer group $\mathcal{L}(s\mathbf{d})$
(001)	$\mathbf{a} \quad \mathbf{b} \quad \mathbf{c}$	$P4_1$	$[s\mathbf{d}, (s + \frac{1}{4})\mathbf{d},$ $(s + \frac{1}{2})\mathbf{d}, (s + \frac{3}{4})\mathbf{d}]$	$p1$ L01

No. 77 $P4_2$

$$\mathcal{G} = P4_2$$

 C_4^3

Orientation orbit (hkl)	Conventional basis of the scanning group $\mathbf{a}' \quad \mathbf{b}' \quad \mathbf{d}$	Scanning group \mathcal{H}	Linear orbit $s\mathbf{d}$	Sectional layer group $\mathcal{L}(s\mathbf{d})$
(001)	$\mathbf{a} \quad \mathbf{b} \quad \mathbf{c}$	$P4_2$	$[s\mathbf{d}, (s + \frac{1}{2})\mathbf{d}]$	$p112$ L03

No. 78 $P4_3$

$$\mathcal{G} = P4_3$$

 C_4^4

Orientation orbit (hkl)	Conventional basis of the scanning group $\mathbf{a}' \quad \mathbf{b}' \quad \mathbf{d}$	Scanning group \mathcal{H}	Linear orbit $s\mathbf{d}$	Sectional layer group $\mathcal{L}(s\mathbf{d})$
(001)	$\mathbf{a} \quad \mathbf{b} \quad \mathbf{c}$	$P4_3$	$[s\mathbf{d}, (s + \frac{1}{4})\mathbf{d},$ $(s + \frac{1}{2})\mathbf{d}, (s + \frac{3}{4})\mathbf{d}]$	$p1$ L01

No. 88 $I4_1/a$ $\mathcal{G} = I4_1/a$ origin 2 C_{4h}^6

Orientation orbit (hkl)	Conventional basis of the scanning group $\mathbf{a}' \quad \mathbf{b}' \quad \mathbf{d}$	Scanning group \mathcal{H}	Linear orbit $s\mathbf{d}$	Sectional layer group $\mathcal{L}(s\mathbf{d})$
(001)	$\mathbf{a} \quad \mathbf{b} \quad \mathbf{c}$	$I4_1/a$ (origin 2)	$[0\mathbf{d}, \frac{1}{2}\mathbf{d};$ $\frac{1}{4}\mathbf{d}, \frac{3}{4}\mathbf{d}]$ $[\frac{1}{8}\mathbf{d}, \frac{5}{8}\mathbf{d};$ $\frac{3}{8}\mathbf{d}, \frac{7}{8}\mathbf{d}]$ $[\pm s\mathbf{d}, (\pm s + \frac{1}{4})\mathbf{d},$ $(\pm s + \frac{1}{2})\mathbf{d}, (\pm s + \frac{3}{4})\mathbf{d}]$	$p112/b$ L07 $p112/a [(\mathbf{a} + \mathbf{b})/4]$ L07 $p\bar{4} (\mathbf{b}/4)$ L50 $p\bar{4} (3\mathbf{b}/4)$ L50 $p112 (\mathbf{b}/4)$ L03

Auxiliary tables for Laue class $C_{4h} - 4/m$ Centring types P and I

Orientation orbit (hkl)	Conventional basis of the scanning group $\mathbf{a}' \quad \mathbf{b}' \quad \mathbf{d}$	Auxiliary basis of the scanning group $\hat{\mathbf{a}} \quad \hat{\mathbf{b}} \quad \hat{\mathbf{c}}$
($mn0$)	$\mathbf{c} \quad n\mathbf{a} - m\mathbf{b} \quad p\mathbf{a} + q\mathbf{b}$	$\mathbf{a} \quad \mathbf{b} \quad \mathbf{c}$
($\bar{m}n0$)	$\mathbf{c} \quad m\mathbf{a} + n\mathbf{b} \quad -q\mathbf{a} + p\mathbf{b}$	

Arithmetic classes $4P$ and $4I$

Serial No.	75	76	77	78	79	80
Group type	C_4^1	C_4^2	C_4^3	C_4^4	C_4^5	C_4^6
Group	$P4$	$P4_1$	$P4_2$	$P4_3$	$I4$	$I4_1$
($mn0$)	$P112$	$P112_1$	$P112$	$P112_1$	$I112$	$I112$
($\bar{m}n0$)						

Arithmetic classes $\bar{4}P$ and $\bar{4}$

Serial No.	81	82
Group type	S_4^1	S_4^2
Group	$P\bar{4}$	$I\bar{4}$
($mn0$)	$P112$	$I112$
($\bar{m}n0$)		

Arithmetic class $4/mP$

Serial No. Group type Group	83 C_{4h}^1 $P4/m$	84 C_{4h}^2 $P4_2/m$	85 C_{4h}^3 $P4/n$		86 C_{4h}^4 $P4_2/n$	
			Origin 1	Origin 2	Origin 1	Origin 2
($mn0$)	$P112/m$	$P112/m$	$P112/n$	$P112/n$	$P112/n$	$P112/n$
($\bar{m}n0$)			$(\mathbf{a} + \mathbf{b})/4$		$(\mathbf{a} + \mathbf{b} + \mathbf{c})/4$	