

13.4. NONCRYSTALLOGRAPHIC SYMMETRY AVERAGING




Fig. 13.4.8.1. Interpolation box for finding the approximate electron density at $G(\Delta x, \Delta y, \Delta z)$, given the eight densities at the corners of the box. The interpolated value can be built up by first using interpolations to determine the densities at A, B, C and D . A second linear interpolation then determines the density at E (from densities at A and B) and at F (from densities at C and D). The third linear interpolation determines the density at G from the densities at E and F . [Reproduced with permission from Rossmann *et al.* (1992). Copyright (1992) International Union of Crystallography.]

13.4.8. Interpolation

Some thought must go into defining the size of the grid interval. Shannon's sampling theorem shows that the grid interval must never be greater than half the limiting resolution of the data. Thus, for instance, if the limiting resolution is 3 Å, the grid intervals must be smaller than 1.5 Å. Clearly, the finer the grid interval, the more accurate the interpolated density, but the computing time will increase with the inverse cube of the size of the grid step. Similarly, if the grid interval is fine, less care and fewer points can be used for interpolation, thus balancing the effect of the finer grid in terms of computing time. In practice, it has been found that an eight-point interpolation (as described below) can be used, provided the grid interval is less than 1/2.5 of the resolution (Rossmann *et al.*, 1992). Other interpolation schemes have also been used (*e.g.* Bricogne, 1976; Nordman, 1980; Hogle *et al.*, 1985; Bolin *et al.*, 1993).

A straightforward 'linear' interpolation can be discussed with reference to Fig. 13.4.8.1 (in mathematical literature, this is called a trilinear approximation or a tensor product of three one-dimensional linear interpolants). Let G be the position at which the density is to be interpolated, and let this point have the fractional grid coordinates $\Delta x, \Delta y, \Delta z$ within the box of surrounding grid points. Let 000 be the point at $\Delta x = 0, \Delta y = 0, \Delta z = 0$. Other grid points will then be at 100, 010, 001 *etc.*, with the point diagonally opposite the origin at 111.

The density at A (between 000 and 100) can then be approximated as the value of the linear interpolant of ρ_{000} and ρ_{100} :

$$\rho(A) \cong \rho_A = \rho_{000} + (\rho_{100} - \rho_{000})\Delta x.$$

Similar expressions for $\rho(B)$, $\rho(C)$ and $\rho(D)$ can also be written. Then, it is possible to calculate an approximate density at E from

$$\rho(E) \cong \rho_E = \rho_A + (\rho_B - \rho_A)\Delta y,$$

with a similar expression for $\rho(F)$. Finally, the interpolated density at G between E and F is given by

$$\rho(G) \cong \rho_G = \rho_E + (\rho_F - \rho_E)\Delta z.$$

Putting all these together, it is easy to show that

$$\begin{aligned} \rho_G = & \rho_{000} + \Delta x(\rho_{100} - \rho_{000}) + \Delta y(\rho_{010} - \rho_{000}) + \Delta z(\rho_{001} - \rho_{000}) \\ & + \Delta x\Delta y(\rho_{000} + \rho_{110} - \rho_{100} - \rho_{010}) \\ & + \Delta y\Delta z(\rho_{000} + \rho_{011} - \rho_{010} - \rho_{001}) \\ & + \Delta z\Delta x(\rho_{000} + \rho_{101} - \rho_{001} - \rho_{100}) \\ & + \Delta x\Delta y\Delta z(\rho_{100} + \rho_{010} + \rho_{001} + \rho_{111} - \rho_{000} - \rho_{101} \\ & - \rho_{011} - \rho_{110}). \end{aligned}$$

13.4.9. Combining different crystal forms

Frequently, a molecule crystallizes in a variety of different crystal forms [*e.g.* hexokinase (Fletterick & Steitz, 1976), the influenza virus neuraminidase spike (Varghese *et al.*, 1983), the histocompatibility antigen HLA (Bjorkman *et al.*, 1987) and the CD4 receptor (Wang *et al.*, 1990)]. It is then advantageous to average between the different crystal forms. This can be achieved by averaging each crystal form independently into a standard orientation in the h -cell (if the redundancy is $N = 1$ for a given crystal form, then this simply amounts to producing a skewed representation of the p -cell in the h -cell environment). The different results, now all in the same h -cell orientation, can be averaged. However, care must be taken to put equal weight on each molecular copy. If the i th cell contains N_i noncrystallographic copies, then the average of the densities, $\rho_i(\mathbf{x})$ ($i = 1, 2, \dots, I$), is

$$\frac{\sum_i N_i \rho_i(\mathbf{x})}{\sum_i N_i}$$

at each grid point, \mathbf{x} , in the h -cell. Additional weights can be added to account for the subjective assessment of the quality of the electron densities in the different crystal cells.

With the h -cell density improved by averaging among different crystal forms, it can now be replaced into the different p -cells. These p -cells can then be back-transformed in the usual manner to obtain a better set of phases. These, in turn, can be associated with the observed structure amplitudes for each p -cell structure, and the cycle can be repeated.

13.4.10. Phase extension and refinement of the NCS parameters

Fourier back-transformation of the modified (averaged and solvent-flattened) map leads to poor phase information immediately outside the previously used resolution limit. If no density modification had been made, the Fourier transform would have yielded exactly the same structure factors as had been used for the original map. However, the modifications result in small structure amplitudes just beyond the previous resolution limit. The resultant phases can then be used in combination with the observed amplitudes in the next map calculation, thus extending the limit of resolution.

If the cell edge of an approximately cubic unit cell is a , and the approximate radius of the molecule is \mathcal{R} (therefore, $\mathcal{R} < a$), then the first node of a spherical diffraction function will occur when $HR = 0.7$, where H is the length of the reciprocal-lattice vector between the closest previously known structure factor and the structure factor just outside the resolution limit. Let $H = n(1/a)$, and let it be assumed that the diffraction-function amplitude is negligible when $HR > 0.7$. Thus, for successful extension, $n = a/\mathcal{R}$. In general, that means that phase extension should be less than two reciprocal-lattice units in one step.

As phase extension proceeds, the accuracy of the NCS elements and the boundaries of the envelope must be constantly improved and updated to match the improved resolution. Arnold & Rossmann