

REFERENCES

- Zhou, Z. H., Hardt, S., Wang, B., Sherman, M. B., Jakana, J. & Chiu, W. (1996). CTF determination of images of ice-embedded single particles using a graphics interface. *J. Struct. Biol.* **116**, 216–222.
- Zhou, Z. H., Macnab, S. J., Jakana, J., Scott, L. R., Chiu, W. & Rixon, F. J. (1998). Identification of the sites of interaction between the scaffold and outer shell in herpes simplex virus-1 capsids by difference electron imaging. *Proc. Natl Acad. Sci. USA*, **95**, 2778–2783.
- Zhou, Z. H., Prasad, B. V. V., Jakana, J., Rixon, F. J. & Chiu, W. (1994). Protein subunit structures in the herpes simplex virus A-capsid determined from 400 kV spot-scan electron cryomicroscopy. *J. Mol. Biol.* **242**, 456–469.
- Zhu, J., Penczek, P. A., Schroder, R. & Frank, J. (1997). Three-dimensional reconstruction with contrast transfer function correction from energy-filtered cryoelectron micrographs: procedure and application to the 70S *Escherichia coli* ribosome. *J. Struct. Biol.* **118**, 197–219.
- 19.7**
- Banci, L., Bertini, I., Cremonini, M. A., Gori-Savellini, G., Luchinat, C., Wüthrich, K. & Güntert, P. (1998). PSEUDYANA for NMR structure calculation of paramagnetic metalloproteins using torsion angle molecular dynamics. *J. Biomol. NMR*, **12**, 553–557.
- Bax, A. & Grzesiek, S. (1993). Methodological advances in protein NMR. *Acc. Chem. Res.* **26**, 131–138.
- Billeter, M. (1992). Comparison of protein structures determined by NMR in solution and by X-ray diffraction in single-crystals. *Q. Rev. Biophys.* **25**, 325–377.
- Billeter, M., Braun, W. & Wüthrich, K. (1982). Sequential resonance assignments in protein ¹H nuclear magnetic resonance spectra: computation of sterically allowed proton–proton distances and statistical analysis of proton–proton distances in single crystal protein conformations. *J. Mol. Biol.* **155**, 321–346.
- Billeter, M., Güntert, P., Luginbühl, P. & Wüthrich, K. (1996). Hydration and DNA recognition by homeodomains. *Cell*, **85**, 1057–1065.
- Braun, W., Epp, O., Wüthrich, K. & Huber, R. (1989). Solution of the phase problem in the X-ray diffraction method for proteins with the nuclear magnetic resonance solution structure as initial model. *J. Mol. Biol.* **206**, 669–676.
- Brünger, A. T., Clore, G. M., Gronenborn, A. M. & Karplus, M. (1986). Three-dimensional structure of proteins determined by molecular dynamics with interproton distance restraints. Application to crambin. *Proc. Natl Acad. Sci. USA*, **83**, 3801–3805.
- Brunne, R. M., Liepinsh, E., Otting, G., Wüthrich, K. & van Gunsteren, W. F. (1993). Hydration of proteins. A comparison of experimental residence times of water molecules solvating the bovine pancreatic trypsin inhibitor with theoretical model calculations. *J. Mol. Biol.* **231**, 1040–1048.
- Brüschweiler, R. & Wright, P. E. (1994). Water self-diffusion model for protein–water NMR cross relaxation. *Chem. Phys. Lett.* **229**, 75–81.
- Cavanagh, J., Fairbrother, W. J., Palmer, A. G. III & Skelton, N. J. (1996). *Protein NMR spectroscopy, principles and practice*. New York: Academic Press.
- Denisov, V. P., Peters, J., Hörlein, H. D. & Halle, B. (1996). Using buried water molecules to explore the energy landscape of proteins. *Nature Struct. Biol.* **3**, 505–509.
- Dubs, A., Wagner, G. & Wüthrich, K. (1979). Individual assignments of amide proton resonances in the proton NMR spectrum of the basic pancreatic trypsin inhibitor. *Biochim. Biophys. Acta*, **577**, 177–194.
- Ernst, R. R., Bodenhausen, G. & Wokaun, A. (1987). *Principles of nuclear magnetic resonance in one and two dimensions*. Oxford: Clarendon Press.
- Güntert, P., Berndt, K. D. & Wüthrich, K. (1993). The program ASNO for computer-supported collection of NOE upper distance constraints as input for protein structure determination. *J. Biomol. NMR*, **3**, 601–606.
- Güntert, P., Mumenthaler, C. & Wüthrich, K. (1997). Torsion angle dynamics for NMR structure calculation with the new program DYANA. *J. Mol. Biol.* **273**, 283–298.
- Havel, T. F. & Wüthrich, K. (1985). An evaluation of the combined use of nuclear magnetic resonance and distance geometry for the determination of protein conformations in solution. *J. Mol. Biol.* **182**, 281–294.
- Kallen, J., Spitzfaden, C., Zurini, M. G. M., Wider, G., Widmer, H., Wüthrich, K. & Walkinshaw, M. D. (1991). Structure of human cyclophilin and its binding site for cyclosporin A determined by X-ray crystallography and NMR spectroscopy. *Nature (London)*, **353**, 276–279.
- Kay, L. E. & Gardner, K. H. (1997). Solution NMR spectroscopy beyond 25 kDa. *Curr. Opin. Struct. Biol.* **7**, 722–731.
- Otting, G., Liepinsh, E. & Wüthrich, K. (1991). Protein hydration in aqueous solution. *Science*, **254**, 974–980.
- Otting, G., Liepinsh, E. & Wüthrich, K. (1993). Disulfide bond isomerization in BPTI and BPTI(G36S): an NMR study of correlated mobility in proteins. *Biochemistry*, **32**, 3571–3582.
- Peng, J. W. & Wagner, G. (1992). Mapping of spectral density functions using heteronuclear NMR relaxation measurements. *J. Magn. Reson.* **98**, 308–332.
- Pervushin, K., Riek, R., Wider, G. & Wüthrich, K. (1997). Attenuated T₂ relaxation by mutual cancellation of dipole–dipole coupling and chemical shift anisotropy indicates an avenue to NMR structures of very large biological macromolecules in solution. *Proc. Natl Acad. Sci. USA*, **94**, 12366–12371.
- Qian, Y. Q., Billeter, M., Otting, G., Müller, M., Gehring, W. J. & Wüthrich, K. (1989). The structure of the Antennapedia homeodomain determined by NMR spectroscopy in solution: comparison with prokaryotic repressors. *Cell*, **59**, 573–580.
- Riek, R., Wider, G., Pervushin, K. & Wüthrich, K. (1999). Polarization transfer by cross-correlated relaxation in solution NMR with very large molecules. *Proc. Natl Acad. Sci. USA*, **96**, 4918–4923.
- Salzmann, M., Pervushin, K., Wider, G., Senn, H. & Wüthrich, K. (1998). TROSY in triple-resonance experiments: new perspectives for sequential NMR assignment of large proteins. *Proc. Natl Acad. Sci. USA*, **95**, 13585–13590.
- Tjandra, N. & Bax, A. (1997). Direct measurement of distances and angles in biomolecules by NMR in dilute liquid crystalline medium. *Science*, **278**, 1111–1114.
- Wagner, G. (1980). Activation volumes for the rotational motion of interior aromatic rings in globular proteins determined by high resolution ¹H NMR at variable pressure. *FEBS Lett.* **112**, 280–284.
- Wagner, G. & Wüthrich, K. (1982). Sequential resonance assignments in protein ¹H nuclear magnetic resonance spectra: basic pancreatic trypsin inhibitor. *J. Mol. Biol.* **155**, 347–366.
- Wishart, D. S., Sykes, B. D. & Richards, F. M. (1991). Relationship between nuclear-magnetic-resonance chemical shift and protein secondary structure. *J. Mol. Biol.* **222**, 311–333.
- Wüthrich, K. (1976). *NMR in biological research: peptides and proteins*. Amsterdam: North Holland.
- Wüthrich, K. (1986). *NMR of proteins and nucleic acids*. New York: Wiley.
- Wüthrich, K. (1989). Protein structure determination in solution by nuclear magnetic resonance spectroscopy. *Science*, **243**, 45–50.
- Wüthrich, K. (1995). NMR – this other method for protein and nucleic acid structure determination. *Acta Cryst.* **D51**, 249–270.
- Wüthrich, K., Billeter, M. & Braun, W. (1984). Polypeptide secondary structure determination by nuclear magnetic resonance observation of short proton–proton distances. *J. Mol. Biol.* **180**, 715–740.
- Wüthrich, K. & Wagner, G. (1975). NMR investigations of the dynamics of the aromatic amino acid residues in the basic pancreatic trypsin inhibitor. *FEBS Lett.* **50**, 265–268.