

## 23.2. Protein–ligand interactions

BY A. E. HODEL AND F. A. QUIOCHO

### 23.2.1. Introduction

There are currently over a thousand unique protein-bound ligands described in the Protein Data Bank, illustrating the enormous variety of small molecules with which proteins interact. These ligands serve as cofactors in protein-mediated reactions, substrates in these reactions, and elements that maintain or alter protein structure or macromolecular assembly. The specific binding of small molecules to proteins is a primary means by which living systems interact and exchange information with their environment. The atomic details of protein–ligand interactions are often quite similar to the intramolecular interactions observed within a protein molecule. Examples of all the various non-covalent interactions described in Parts 20 and 22, such as hydrogen bonds, van der Waals forces and other electrostatic phenomena, are observed between proteins and their small-molecule ligands.

Through the diverse interactions observed between proteins and their ligands, a few fundamental patterns of recognition emerge. In general, ligand binding requires that the protein partially or fully sequesters the ligand from the solvent. This demands that the energy of interaction between the protein and the ligand must be strong enough to overcome the interactions between both species and the solvent as well as the translational and rotational entropy which is lost upon fixing the orientation of the ligand relative to the protein. The protein achieves this level of interaction by presenting a binding site that is complementary to the ligand both in shape and electrostatic functionality. Beyond this generalization, each ligand has its own unique and complicated story. Rather than attempt to summarize the enormous subject of protein–ligand interactions in a comprehensive manner, we will instead illustrate several of the unique interactions observed between proteins and other molecules.

### 23.2.2. Protein–carbohydrate interactions

The interaction between proteins and carbohydrates provides a prototypic example of how proteins specifically recognize small organic ligands. Protein-mediated recognition of carbohydrates is crucial in a diverse array of processes, including the transport, biosynthesis and storage of carbohydrates as an energy source, signal transduction through carbohydrate messengers, and cell–cell recognition and adhesion (Rademacher *et al.*, 1988). Many aspects of protein–carbohydrate recognition are observed in the interactions of proteins with other small-molecule ligands. For example, carbohydrate recognition is largely conferred through a combination of hydrogen bonding and van der Waals interactions with the protein (Quiucho, 1986). These interactions are generally presented in a binding site that is highly complementary to the target ligand in shape and functionality. Carbohydrate-binding proteins also occasionally employ ordered water molecules and bound metal ions to facilitate ligand recognition (Quiucho *et al.*, 1989; Vyas, 1991; Weis & Drickamer, 1996). There are also many examples of ‘induced fit’ recognition of carbohydrates, where the binding of the ligand induces a conformational change in the protein. This phenomenon of ligand-induced conformational change was observed in one of the first ligand–protein interactions visualized by X-ray crystallography, namely the binding of a carbohydrate substrate to lysozyme (Blake *et al.*, 1967). Since these early studies, a wide variety of carbohydrate–protein structures have been determined, and a number of general themes have emerged from this continually active field (Vyas, 1991). These themes are further applicable in the general study of protein–ligand recognition.

Two general classes of carbohydrate-binding modes have been observed in their complexes with proteins (Quiucho, 1986; Vyas,

1991). The first group of proteins completely sequester the carbohydrate from the surrounding solvent. These proteins, including the periplasmic proteins and the catalytic site of glycogen phosphorylase, tend to have a high affinity for their ligand ( $K_d \approx 10^{-6}$ – $10^{-7}$ ). The high affinity of these proteins can be attributed to the entropic gain of desolvating the protein and carbohydrate surfaces as well as the exceptionally high degree of functional complementarity in the ligand-binding site. The periplasmic proteins nearly saturate the hydrogen-bonding potential of their carbohydrate ligands (Quiucho, 1986). The second group of proteins bind their ligands in shallow clefts in the solvent-exposed protein surface. This mode of binding, observed in the lectins, lysozyme and the storage site of glycogen phosphorylase, generally has a lower binding constant than the first class of binding proteins ( $K_d \approx 10^{-3}$ – $10^{-6}$ ). Some members of this second class of proteins, such as the lectins, are able to increase the affinity for their ligands dramatically by clustering a number of low-affinity sites through the oligomerization of the polypeptides (Weis & Drickamer, 1996). The specific arrangement of lectin oligomers allows them to discriminate between large cell-surface arrays of polysaccharides with high affinity and selectivity.

#### 23.2.2.1. Carbohydrate recognition at the atomic level

An early review of protein–carbohydrate interactions revealed several atomic level interactions that continue to appear ubiquitously in the structures of protein–carbohydrate complexes (Quiucho, 1986). A generic cyclic sugar in mono- or oligosaccharides appears to be recognized as a disc that displays two flat non-polar surfaces surrounded by a ring of polar hydroxyls. Proteins recognize these features by hydrogen bonding to the ring of polar hydroxyls while stacking flat aromatic side chains against the non-polar disc faces.

Cooperative hydrogen bonding, where the hydroxyl group of a carbohydrate participates as both a donor and acceptor of hydrogen bonds, is often observed in the direct interactions between proteins and carbohydrate ligands (Fig. 23.2.2.1). The  $sp^3$ -hybridized oxygen atom of a carbohydrate hydroxyl may act as both an acceptor of two hydrogen bonds through the two lone pairs of electrons as well as a donor of a single hydrogen bond. Cooperative




Fig. 23.2.2.1. The atomic recognition of carbohydrates by a protein. The environment of glucose bound to the galactose/glucose binding protein is shown (Vyas *et al.*, 1988). (a) The ring of hydrogen bonds around the polar edge of the sugar molecule. Note the ‘bidentate’ hydrogen bonds with Asp236, Arg158 and Asn91. As the hydroxyl groups 1, 2 and 3 simultaneously donate and accept hydrogen bonds from protein side chains, they are involved in ‘cooperative’ hydrogen bonds. (b) Two aromatic protein side chains, Phe16 and Trp183, ‘stack’ on the non-polar faces of the carbohydrate.

## 23. STRUCTURAL ANALYSIS AND CLASSIFICATION

Table 23.2.3.1. *Metal ions associated with proteins*

Metal ion	Concentration in blood plasma (mM)	Common cofactors	Hard/soft classification	Common coordination number and geometry	Preferred ligand atom
Na <sup>+</sup>	138		Hard	6	O
K <sup>+</sup>	4		Hard	8	O
Ca <sup>2+</sup>	3		Hard	8	O
Mg <sup>2+</sup>	1		Hard	6	O
Fe	0.02	Haem	Intermediate	6 (octahedral)	N
Zn <sup>2+</sup>	0.02		Intermediate	4 (tetrahedral), 6 (octahedral)	N, S
Cu <sup>2+</sup>	0.015		Soft	4 (tetrahedral)	S
Co <sup>2+</sup>	0.002		Hard	6 (octahedral)	O
Mn <sup>2+</sup>	0		Hard	6 (octahedral)	O
Ni <sup>2+</sup>	0		Intermediate	6 (octahedral)	N
Mo	0	Pterin	Intermediate	6 (octahedral)	S
W	0	Pterin	Intermediate	6 (octahedral)	S
V	0			5 (trigonal bipyramidal)	

hydrogen bonding generally follows a simple pattern in which the carbohydrate hydroxyl accepts a hydrogen bond from a protein amide group while simultaneously donating a hydrogen bond to a protein carbonyl oxygen. Hydrogen bonding to protein hydroxyl groups is observed only infrequently. This pattern is thought to be a result, in part, of the entropic cost of fixing a freely rotating protein hydroxyl group while simultaneously fixing the ligand hydroxyl group. Amides and carbonyls are usually fixed in a planar geometry and thus do not require as much energy to compensate for their loss of entropy in ligand binding.

The vicinal hydroxyl groups of carbohydrates provide an ideal geometry for the formation of 'bidentate' hydrogen bonds, where the pair of hydroxyls interacts with two functional groups of a single amino-acid side chain or the main-chain amide groups of two consecutive residues (Fig. 23.2.2.1). These interactions occur when the adjacent carbohydrate hydroxyls are either both equatorial, or one is equatorial and the other axial. The interatomic distance for the carbohydrate hydroxyl oxygens is ~2.8 Å in these cases, allowing for a bidentate interaction with the planar side chains of aspartate, asparagine, glutamate, glutamine and arginine. Bidentate hydrogen bonds have not been observed for consecutive axial hydroxyls where the oxygen–oxygen distance increases to ~3.7 Å.

Carbohydrates often present a disc-like face of non-polar aliphatic hydrogen atoms which proteins recognize through the use of aromatic side chains. The protein aromatic groups are 'stacked' on the flat face of the carbohydrate, thus generating both specificity and binding energy through van der Waals interactions. Tryptophan, the aromatic amino acid with the largest surface area and highest electronegativity, is the most common side chain employed in van der Waals 'stacking' with carbohydrates. The infrequent use of aliphatic groups in the binding of the non-polar carbohydrate faces suggests that the aromatic moieties are employed in a specific manner. The electron-rich electron clouds of the aromatic side chains may provide a strong electrostatic interaction with the aliphatic carbohydrate protons that could not be satisfied by protein aliphatic groups. The anionic character of aromatic side chains is observed in a number of protein–intramolecular (Chapter 22.2) and protein–ligand interactions (see below).

### 23.2.3. Metals

Metal ions provide a number of important functions in their diverse and ubiquitous interactions with proteins. The most common

function for a protein-bound metal ion is the stabilization and orientation of the protein tertiary structure through coordination to specific protein functional groups. In addition to this structural role, metal ions are also often directly involved in enzyme catalysis and protein function. Examples of these functions include redox reactions, the activation of chemical bonds and the binding of specific ligands. Myoglobin, the first protein structure determined by X-ray crystallography, specifically binds molecular oxygen through an iron ion of a haem cofactor. Myoglobin provides a prototypic example of a protein and a metal ion providing a unique and specific functionality through their combination.

#### 23.2.3.1. *Metals important in protein function and structure*

A number of metals are relatively abundant and available in living systems (Table 23.2.3.1) (Glusker, 1991). The most common ions include sodium, potassium, magnesium and calcium. Along with these ions, a large variety of trace metals are also found coordinated to proteins. The structures of protein complexes with some of these trace ions, including iron, zinc and copper, have been studied extensively for some time (Glusker, 1991). More recently, the structures of protein complexes with more unusual ions, such as nickel, vanadium and tungsten, have been determined (Volbeda *et al.*, 1996).

Specificity in the interactions between proteins and metal ions is conferred through each ion's preference for the coordinating atoms and the geometry of the binding site. All four of the more common metals, *i.e.* sodium, potassium, magnesium and calcium, are classified as 'hard' metals, referring to the polarizability of the electron cloud of the ion. The nucleus of a hard metal has a relatively tight hold on the surrounding electrons. These ions lack easily excitable unshared electrons and have a low polarizability. The interactions between these metals and their ligands tend to have the character of ionic interactions rather than the more covalent nature preferred by the 'soft' metals. In general, the hard metals prefer to coordinate with hard acids, such as the oxygen atoms of hydroxyls, carbonyls and carboxyls.

The soft metals have a high polarizability, large ionic radius and several unshared valence electrons. They generally prefer to coordinate with soft acids, such as the thiol and thiol ether groups of cysteine and methionine. The loosely held valence electrons of soft metals tend to favour partially covalent  $\pi$ -bonding with their coordinated ligands. These outer-shell electrons can be donated to the empty outer orbitals of the ligand atom. The partially covalent nature of these bonds yields more stable complexes than the ionic