

23. STRUCTURAL ANALYSIS AND CLASSIFICATION




Fig. 23.4.3.5. Schematic illustration of water molecules bound in different types of grooves between protein and ligand. The hatched surfaces represent the ligand surface. (a) Water molecules bound in an indentation on the protein surface, where the protein surface area exposed to the water molecules is far larger than the ligand surface area; (b) water molecules bound in indentations on the ligand surface, where the ligand surface area exposed to the water molecule is larger than the protein surface area; (c) water molecules bound in shallow grooves at the protein–ligand interface and on the ligand surface; and (d) water molecules bound in clusters in elongated grooves with micro-grooves. Reprinted with permission from Poormina & Dean (1995c). Copyright (1995) Kluwer Academic Publishers.

bridging water molecules in deep grooves on the protein or on the ligand, respectively. The most common situation is illustrated in Fig. 23.4.3.5(a), with that in Fig. 23.4.3.5(b) occurring very rarely. Fig. 23.4.3.5(c) shows the situation where water molecules are found to interact with the ligand alone or at the periphery of the protein–ligand interface. Finally, Fig. 23.4.3.5(d) illustrates the situation where clusters of water molecules occupy elongated grooves, mediating the protein–ligand interaction. A striking example of this is given by the complex between chloramphenicol acetyl transferase and chloramphenicol, where two clusters of water molecules are found to form a layer between the enzyme and the ligand (Poormina & Dean, 1995c).

For the purposes of analysis, the authors distinguish between water molecules that interact with both protein and ligand, forming a bridge between the two, and water molecules that interact with either the protein or the ligand, but not with both. There is also a group of water molecules that interact with neither protein nor ligand, but are thought to contribute to the stability of the network of water molecules at the protein–ligand interface.

Of the 58 water molecules found to bridge between protein and ligand, 38 (nearly 80%) make three or more hydrogen bonds and satisfy tetrahedral geometry. Furthermore, they bind in deep grooves, generally interacting more strongly with the protein (Fig. 23.4.3.5a). The *B* factors of these bridging water molecules are comparable to those of the protein atoms with which they interact. They can, in effect, be considered an integral part of the protein structure and binding site. Many of these bridging water molecules are conserved throughout homologous proteins, even when different ligands are considered, and are clearly structurally significant in maintaining the properties of the protein binding sites.

Water molecules found to bind in shallow grooves do so either at the ligand surface or at the periphery of the protein–ligand interface.

For many of these water molecules, the surface areas of the protein and the ligand exposed to the same water molecule are nearly equal. Water molecules binding in shallow grooves are found to have zero to two polar contacts with the protein and are not particularly well conserved within families of homologous proteins.

In general, the authors conclude that water molecules that are to be considered as part of the protein binding site during the design of a new ligand are those that bind in deep grooves, making multiple hydrogen bonds to protein atoms. These water molecules tend to be conserved through families of homologous proteins. The amino-acid residues that interact with deep-groove water molecules tend to be more conserved compared with other residues interacting with the ligand. Conversely, the binding of water in shallow grooves does not seem to be influenced by any special general feature of the protein or ligand surface, and it would be difficult to select water molecules *a priori* for inclusion as part of the protein structure during the process of ligand design.

23.4.4. Water structure in groups of well studied proteins

The analysis of general features of protein–water interactions derived from large databases provides an important context for the study of solvent structure in individual proteins. The number of crystallographically visible water molecules in any one X-ray structure depends on the resolution of the data, the degree of refinement of the model, the criteria used for placement of the less well defined water molecules, and on the experience of the crystallographer. Therefore, to differentiate between water molecules that have functional roles and those that associate randomly with the protein, it is desirable to determine commonalities between several independently solved structures of the protein of interest. There are different types of functional roles that can be determined at several levels. At the global level, one can find a small number of water molecules that are essential for the structural architecture common to a given family of homologous proteins. There are also those water molecules that are structurally important for a specific protein, being present in all independently solved structures of that protein, regardless of the crystal form in which the water molecule was determined or of its interactions with ligands. Water molecules that consistently appear in crystal structures of the protein solved in a specific space group but in no others may be important for crystal packing, but not to the integrity of the protein itself. Finally, a given water molecule may be essential for mediating in a protein–ligand complex, but never appear in the native protein. At this level, all of the independently solved structures of the complex would have the water molecule present. In the examples that follow, comparative analysis between carefully selected groups of structures reveals conserved water molecules at all of these different levels and shows how they carry out particular functional roles in specific examples.

23.4.4.1. Crystal structures of homologous proteins

There are two families of homologous proteins for which extensive solvent-structure comparisons have revealed water molecules important in maintaining structural features common to all members of the family. In the first study presented here, 35 crystal structures of eight members of the serine protease family were analysed (Sreenivasan & Axelsen, 1992), while the second study comprises a similar analysis of 11 independently solved structures of six members of the legume lectin family (Loris *et al.*, 1994).

23.4.4.1.1. Serine proteases of the trypsin family

The serine proteases have an especially large number of buried water molecules. Using a probe sphere of radius 1.4 Å, an iterative