

## 23. STRUCTURAL ANALYSIS AND CLASSIFICATION




Fig. 23.4.4.10. Three-dimensional structure of RNase T1. Secondary structure is denoted as follows:  $\alpha_1$ ,  $\alpha$ -helix;  $\beta_n$ , strands of  $\beta$ -sheet structure;  $L_n$ , loops. Drawn using *MOLSCRIPT* (Kraulis, 1991). Residue numbers indicate the beginning and end of secondary-structure elements. Reprinted with permission from Pletinckx *et al.* (1994). Copyright (1994) American Chemical Society.

guanine. Therefore, the information obtained from a study of the solvent structure in ribonuclease A is completely independent from that described above for ribonuclease T1. A collection of ten crystal structures of ribonuclease A, derived from five different crystal forms, were compared pairwise after least-squares superposition (Zegers *et al.*, 1994). 17 conserved water molecules were found to be within a sphere of 0.5 Å of each other in all of the ten structures




Fig. 23.4.4.11. Overall structure of RNase A. The overall structure of the d(CpA) complex of RNase A is shown as a ribbon drawing using *MOLSCRIPT* (Kraulis, 1991). The conserved water molecules are shown as white spheres and the d(CpA) inhibitor in black. The three helices are labelled H1, H2 and H3. Reprinted with the permission of Cambridge University Press from Zegers *et al.* (1994). Copyright (1994) The Protein Society.

and are shown in Fig. 23.4.4.11. These water molecules were found in small clusters of two or three or as part of a larger solvent network. Not surprisingly, they form multiple hydrogen bonds with the protein and generally have low temperature factors. Of the 17 structurally conserved sites, 13 are associated with one of the three  $\alpha$ -helices. Most of these link the helices to one of the  $\beta$ -strands. Three water molecules are involved in hydrogen bonding with unpaired amido and carbonyl groups on the protein, and one is found on top of the  $\beta$ -pleated sheet. These interactions result in bringing together elements of secondary structure and in stabilizing distortions within these elements. Conserved water molecules are also responsible for bridging the N-terminal helix to the C-terminal  $\beta$ -strand, which form the two halves of the active site.

## 23.4.4.2.5. Protein kinase A

The comparative study of water molecules in seven different protein kinase A structures in complex with different ligands focused exclusively on the active site (Shaltiel *et al.*, 1998). All of the structures were solved from isomorphous crystals to resolutions ranging from 2.0 to 2.9 Å. The more lenient cutoff of 1.5 Å for the radius of the sphere within which the conserved water molecules must be found among the different structures is consistent with the relatively low resolutions of the structures included in this study. The group of structures represents the open, the closed and an intermediate conformation of the catalytic kinase domain. There is a set of six conserved water sites in the active site, in addition to the ATP molecule and the magnesium ion. The conserved water molecules coordinate to ATP, the metal ion and a conserved Tyr residue from the carboxyl terminus of the protein. Thus, the active site consists of an extended network of interactions that weave together both domains of the core, with water molecules playing an integral role in maintaining the structural features important for catalysis. Many of these water molecules associate directly with the inhibitors. In addition, five water sites are observed in positions that would be occupied by substrates or substrate analogues. These water molecules are displaced by ligand oxygen atoms that can compensate for the water hydrogen-bonding interaction with the protein.

## 23.4.4.3. Summary

Water molecules associated with proteins can be divided between those that are conserved as a result of their functional significance and those that are partially conserved or not conserved at all. The conserved water molecules are generally classified as buried or channel (by a variety of criteria). They tend to be present in the clefts between domains, are critical components of active sites, or bridge between secondary-structure elements. The water molecules that are not conserved occupy hydration sites with favourable hydrogen-bonding characteristics, where the presence of a water molecule is not essential for the structural or functional integrity of the protein.

The displacement of water molecules by organic solvent molecules in the elastase work described above showed that most displaced waters are those classified as surface or crystal-contact waters (Mattos *et al.*, 2000). In the three cases where a buried water molecule was displaced, an alcohol hydroxyl oxygen was found to replace the protein–water hydrogen-bonding interactions. This is analogous to the active-site water molecule in the HIV aspartate protease that gets replaced by a carbonyl group of a potent cyclic urea inhibitor (Lam *et al.*, 1994). In these situations, release of a tightly bound water molecule is entropically favourable, and its enthalpic interactions with the protein are compensated by similar protein–ligand interactions.

The effect of crystal contacts on the water structure was clearly illustrated in the T4 lysozyme work (Zhang & Matthews, 1994).