

23. STRUCTURAL ANALYSIS AND CLASSIFICATION




Fig. 23.4.4.3. A $2F_o - F_c$ electron-density map contoured at the 1.2σ level shows a distinct ellipsoidal density for acetonitrile 707 and a spherical density for a nearby water molecule. The protein backbone of the binding pocket is represented with nitrogen atoms shown in dark grey, oxygen atoms in medium grey and carbons in a lighter grey. *MOLSCRIPT* (Kraulis, 1991) was used in the preparation of this figure. Reprinted with permission from Allen *et al.* (1996). Copyright (1996) American Chemical Society.

23.4.4.2.1. Elastase

The crystal structure of porcine pancreatic elastase was solved in a variety of organic solvents, with the primary goal of mapping binding sites on the protein that could accommodate molecules representative of functional groups likely to be found in larger ligands (Ringe, 1995; Mattos & Ringe, 1996; Mattos *et al.*, 2000). Crystals of elastase cross-linked with glutaraldehyde were transferred to the following solutions: 100% acetonitrile, 95% acetone, 55% dimethylformamide, 80% ethanol, 40% trifluoroethanol, 80% isopropanol and 80% 5-hexene-1,2-diol (Allen *et al.*, 1996; Mattos & Ringe, 1996). In general, the crystals did not diffract in most neat organic solvents. However, in the acetonitrile case, where they did, the result was striking. In the structure of elastase solved in >99% acetonitrile, there were 126 water molecules visible in the electron-density maps, indicating that a good portion of the first hydration shell of the protein was still present. In contrast, only nine molecules of acetonitrile were clearly identified in the electron-density maps (Allen *et al.*, 1996). This is a powerful assertion of the evolutionary specificity of water molecules for protein surfaces. Fig. 23.4.4.3 shows the clear contrast between the elongated electron density of an acetonitrile molecule and the spherical electron density of a water molecule.

A similar result was obtained for all of the elastase structures solved in the mixtures of organic solvent and water mentioned above. A total of 11 structures were analysed, each containing 126–177 water molecules. The structures are listed in Table 23.4.4.1, together with the resolution of the data collected, the number of water molecules present and the number of organic solvent molecules observed in each case. The $C\alpha$ superposition of the protein atoms in the 11 structures yielded a total of 1661 individual water molecules, occupying 426 unique water-binding sites on the elastase surface. Given that elastase has a total of 240 amino-acid

residues, this represents a significant portion of the first hydration shell of the protein. This group of elastase structures served as a powerful source of information, leading to a classification of water types according to their interaction with the protein and an analysis of the specificity for water within each of the types determined (Bellamacina *et al.*, 1999).

All of the 1661 water molecules were renumbered according to the site on the protein where they were found. Any two water molecules within 1 Å of a water molecule in the cross-linked elastase structure solved in distilled water (used as the reference structure) have a common number. 39 of the 426 water-binding sites were occupied in every one of the 11 structures and were considered structurally conserved. Among these are the 16 buried water-binding sites thought to be conserved among all serine proteases (Sreenivasan & Axelsen, 1992). The 26 remaining conserved water molecules are specific to elastase and are not necessarily buried. These water molecules in general tend to have low B factors, but a few have B factors in the 30–35 Å² range and one conserved water molecule has a B factor of 42 Å².

The classification of the water sites as buried, channel, crystal contact or surface was based on the number of hydrogen-bonding interactions that a water molecule at the site could make to the protein and involved no surface-accessibility calculations (Bellamacina *et al.*, 1999). Water molecules were classified as buried if they made at least three good hydrogen-bonding interactions with protein main-chain atoms. A total of 23 buried water sites were identified in this manner, including 13 of the sites classified as buried by Sreenivasan & Axelsen (1992). One of the 16 serine protease conserved water-molecule sites is replaced by a His side chain in elastase (Sreenivasan & Axelsen, 1992). The remaining two serine protease conserved water sites were classified as channel based on the criteria used in the present study (see below). Interestingly, with the exception of these two channel water molecules, all of the buried sites found to be conserved in serine proteases are strictly conserved in all of the 11 structures in Table 23.4.4.1. The two channel water molecules are found in the aqueous structures of elastase, but are virtually absent in elastase transferred to organic solvents.

The water molecules occupying the 23 buried water sites identified in this study are tightly clustered when the protein $C\alpha$ atoms are superimposed by least squares, and the interactions with the protein are conserved from structure to structure. Fig. 23.4.4.4 shows the positions of the buried water-binding sites in elastase. In general, they are found in the cleft between the two domains, in bridging elements of the secondary structure and at the base of water channels. This observation is consistent with the current

Table 23.4.4.1. Multiple-solvent crystal structures of elastase

Structure	Resolution (Å)	No. of water molecules	No. of organic solvent molecules
Cross-linked	1.9	165	0
Acetonitrile	2.2	126	9
Acetone	2.0	126	6
Dimethylformamide	2.0	153	6
Ethanol	2.0	135	12
Trifluoroethanol (1)	1.9	175	4
Trifluoroethanol (2)	1.85	177	3
Isopropanol	2.2	160	4
Benzene	1.9	162	4
Cyclohexane	1.95	135	7
5-Hexene-1,2-diol	2.2	147	5