

23. STRUCTURAL ANALYSIS AND CLASSIFICATION




Fig. 23.4.4.6. Elastase structure represented as in Fig. 23.4.4.4. The crystallographic water molecules involved in crystal contacts in 11 superimposed elastase structures solved in a variety of solvents are shown in green.

isopropylanilide (Mattos *et al.*, 1994) or the trifluoroacetyl-Lys-Pro-*p*-trifluoromethylanilide (Mattos *et al.*, 1995) inhibitors in the structures of their complexes with elastase. These inhibitors span a large area of the active site, including an exosite not occupied by

substrate analogue inhibitors (Mattos *et al.*, 1994, 1995). The water-binding sites in the active site are not very well conserved, with most sites represented in only two to four of the 11 structures. When all of the structures are superimposed, there is at least one water molecule in each of the subsites in the elastase active site. These water molecules are displaced either by inhibitors or by organic solvent molecules in the various structures. It is not surprising that in elastase, a protein with relatively broad substrate specificity, the active site in the uncomplexed native protein is populated by many displaceable surface water molecules. With the exception of a water molecule present in the oxyanion hole, these water molecules tend to make a single hydrogen bond with the protein. This hydrogen-bonding interaction is not generally conserved between different structures where a given site is occupied in multiple structures. The displacement of these water molecules upon ligand binding is entropically favourable, as they are released into bulk solvent, without too much enthalpic cost. This relatively small enthalpic cost can be compensated by the protein–ligand interactions.

Fig. 23.4.4.8 shows all of the 1661 water molecules colour-coded by the various classifications described above. Clearly, the entire surface of the protein is well hydrated. Notice how the yellow channel waters are often followed by a red buried water molecule. In addition, there is often no obvious spatial distinction between molecules categorized as crystal contacts (green) and those categorized as surface (blue).

23.4.4.2.2. *T4 lysozyme*

Over 150 mutants of T4 lysozyme have been studied to date, and, for the majority of these, the crystal structures are available. Although most of the mutant structures crystallize isomorphously to the wild type, many of them provide a view of the molecule in different crystal environments. This collection of structures leads to




Fig. 23.4.4.7. Elastase structure represented as in Fig. 23.4.4.4. The surface crystallographic water molecules found in 11 superimposed elastase structures solved in a variety of solvents are shown in blue.




Fig. 23.4.4.8. Elastase structure represented as in Fig. 23.4.4.4. The 1661 water molecules found in 11 superimposed elastase structures of elastase are colour-coded as in Figs. 23.4.4.4–23.4.4.7.