

## 23.4. SOLVENT STRUCTURE

water activity affects site-specific DNA recognition, with an increase in osmotic pressure leading to a decrease in accuracy of protein–DNA recognition, as observed by DNA cleavage at sites containing an incorrect base pair (Robinson & Siglar, 1993). The results of this study strongly imply a role for one or more water molecules in recognition of specific sequences of DNA. The authors suggest that water mediation may constitute a general motif for sequence-specific DNA recognition by DNA-binding proteins (Robinson & Siglar, 1993).

The role of water molecules as mediators of sequence-specific DNA recognition may be a general motif, but not a necessary one. The solution NMR structure of the complex of erythroid transcription factor GATA-1 with the 16-base-pair DNA fragment GTTGCAGATAAACATT, containing the recognition sequence, shows that the specific interactions between GATA-1 and the major groove of the DNA are dominated by van der Waals interactions hydrophobic in nature (Omichinski *et al.*, 1993). Furthermore, NMR experiments designed to identify the location of water molecules in the complex detected clusters of water molecules bridging the protein to the DNA phosphate backbone, but showed that water was excluded from the hydrophobic interface between the protein and the DNA bases (Clore *et al.*, 1994). Although many of the existing crystal structures of protein–DNA complexes support the general view that water molecules are often integral components of the specific recognition between the protein and the target DNA, this solution structure provides an important example of exclusion of water molecules from the specificity determinants. In the GATA-1–DNA complex, however, water molecules do mediate non-specific binding of the protein to the DNA backbone. It appears, not surprisingly, that water molecules play a variety of roles in the mediation of protein–DNA interactions and that these roles are specific to each particular case.

## 23.4.6.3. Cooperativity in dimeric haemoglobin

The X-ray crystal structures of liganded and unliganded dimeric haemoglobin from *Scapharca inaequalvis* have revealed that water molecules at the dimer interface form an integral part of the cooperativity mechanism in this system (Condon & Royer, 1994; Royer, 1994). The binding of oxygen to one of the monomers causes little rearrangement of quaternary structure. It does, instead, displace the side chain of Phe97 which, in the low-affinity deoxy form, packs in the haem pocket (Royer *et al.*, 1990). Phe97 in the deoxy form lowers the oxygen affinity by restricting movement of the iron atom into the haem plane (Royer, 1994). Upon oxygen binding, Phe97 flips to the dimer interface, removing six out of the 17 water molecules that are found in the deoxy form (Fig. 23.4.6.1). The resultant destabilization of the water clusters found between the two subunits facilitates the flipping of Phe97 in the other subunit, with a concomitant increase in oxygen affinity of the haem in the second subunit (Pardanani *et al.*, 1997; Royer *et al.*, 1997).

In each of the monomeric subunits, Thr72 is positioned to form a hydrogen bond with a water molecule at the periphery of the deoxy dimer interface (not shown in Fig. 23.4.6.1). In effect, this interaction caps the water cluster on either side of the interface, presumably helping to stabilize these well ordered water molecules. The isosteric mutation Thr72 to Val was designed to test the importance of this interaction to the stability of the water cluster in the low-affinity haemoglobin dimer and the resultant effect on ligand affinity and cooperativity (Royer *et al.*, 1996). The crystal structure of the T72V mutant was solved to 1.6 Å resolution. This crystal structure reveals that the only significant difference between the mutant and wild-type proteins is the loss of the two water molecules that directly hydrogen-bond to Thr72 in each of the wild-type subunits. Furthermore, there is a significant increase in both




Fig. 23.4.6.1. *Scapharca* HbI interface water molecules. (a) Deoxy-HbI at 1.6 Å resolution (PDB code 3SDH) and (b) HbI-CO at 1.4 Å resolution (PDB code 4SDH). Included is a ribbon diagram showing the tertiary structure of each subunit, bond representations for the haem group and Phe97 side chain, and spheres representing the approximate van der Waals radii of oxygen atoms for core interface water molecules. Note the cluster of 17 ordered water molecules in the interface of deoxy-HbI for which Phe97 is packed in the haem pocket. Upon ligation, by either CO or O<sub>2</sub>, Phe97 is extruded into the interface and disrupts this water cluster, expelling six water molecules from the interface. These plots were produced with the program *MOLSCRIPT* (Kraulis, 1991). Reprinted with permission from Royer *et al.* (1997). Copyright (1997) The American Society for Biochemistry & Molecular Biology.