

24.5. THE PROTEIN DATA BANK, 1999–

Table 24.5.2.2. Demographics of the released data in the PDB as of 14 September 1999

Experimental technique	Molecule type				
	Proteins, peptides, and viruses	Protein–nucleic acid complexes	Nucleic acids	Carbohydrates and other	Total
X-ray diffraction and other	7946	390	439	14	8789
NMR	1365	53	270	4	1692
Theoretical modelling	202	16	15	0	233
Total	9513	459	724	18	10714

contacts, and positions of water molecules. In consultation with the community, other structure checks will be implemented over the next few years.

The experimental data are also checked. Currently, X-ray crystallographic data are validated and plans for checking NMR data are in progress. For X-ray crystallographic structures, the structure factors are validated using *SFCHECK* (Vaguine *et al.*, 1999). This program extracts the deposited *R* factor, resolution and model information, and then compares them with values calculated from coordinate and structure-factor files. It also calculates an overall *B* factor, coordinate errors, an effective resolution and completeness. The summary of the density correlation shift and *B* factor are reported for each residue. As specific procedures are developed for checking NMR structures against experimental data, they will be incorporated into the PDB validation procedures.

24.5.2.3. NMR data

The PDB staff recognize that NMR data need a special development effort. Historically these data have been retro-fitted into a PDB format defined around crystallographic information. As a first step towards improving this situation, the PDB carried out an extensive assessment of the current NMR holdings and presented the findings to a task force consisting of a cross section of NMR researchers. The PDB is working with this group, the BioMag-ResBank (BMRB; Ulrich *et al.*, 1989) and other members of the NMR community to develop an NMR data dictionary along with deposition and validation tools specific for NMR structures.

24.5.2.4. Data-processing statistics

Production processing of PDB entries by the RCSB began on 27 January 1999. As of 1 July 1999, when the RCSB became fully responsible for the PDB, approximately 80% of all structures submitted to the PDB are deposited *via ADIT* and processed by the RCSB. Another 20% are submitted *via AutoDep* to the European Bioinformatics Institute (EBI), who process these submissions and forward them to the PDB for archiving and distribution. The average time from deposition to the completion of data processing including author interactions is two weeks. The number of structures with a HOLD release status remains at about 20% of all submissions; 57% are held until publication (HPUB); and 23% are released immediately after processing.

Table 24.5.2.2 shows the breakdown of the types of structures in the PDB. As of 14 September 1999, the PDB contained 10 714 publicly accessible structures with another 1169 entries on hold (not shown). Of these, 8789 (82%) were determined by X-ray methods, 1692 (16%) were determined by NMR and 233 (2%) were theoretical models. Overall, 35% of the entries have deposited experimental data.

24.5.3. The PDB database resource

24.5.3.1. The database architecture

In recognition of the fact that no single architecture can fully express the information content of the PDB, an integrated system of heterogeneous databases and indices that store and organize the structural data has been created. At present there are five major components (Fig. 24.5.3.1):

(1) The core relational database managed by Sybase (Sybase Inc., 1995) provides the central physical storage for the primary experimental and coordinate data described in Table 24.5.2.1. The core PDB relational database contains all deposited information in a tabular form that can be accessed across any number of structures.

(2) The final curated data files (in PDB format) and data dictionaries are the archival data and are present as ASCII files in the ftp archive.

(3) The POM-based databases (Shindyalov & Bourne, 1997) consist of indexed objects containing native (*e.g.* atomic coordinates) and derived properties (*e.g.* calculated secondary-structure assignments and property profiles). Some properties require no derivation, for example, *B* factors; others must be derived, for example, exposure of each amino-acid residue (Lee & Richards, 1971) or *C α* contact maps. Properties requiring significant computation time, such as structure neighbours (Shindyalov & Bourne, 1998), are pre-calculated when the database is incremented to save considerable user-access time.

(4) The Biological Macromolecule Crystallization Database (BMCD; Gilliland, 1988) is organized as a relational database within Sybase and contains three general categories of literature-derived information: macromolecular, crystal and summary data.

(5) The Netscape LDAP server is used to index the textual content of the PDB in a structured format and provides support for keyword searches.

In the current implementation, communication among databases has been accomplished using the common gateway interface (CGI). An integrated web interface dispatches a query to the appropriate database(s), which then executes the query. Each database returns the PDB identifiers that satisfy the query, and the CGI program




Fig. 24.5.3.1. The integrated query interface to the PDB.