

## 5.2. CRYSTAL-DENSITY MEASUREMENTS

be calculated in advance, and looked for first. The positions of calibration drops and of crystals in these centrifuge tubes can be measured with a hand-held ruler to a resolution of about 0.5 mm.

Particularly for crystals with high values of  $V_M$  (i.e., loosely packed) or for crystals of large molecular weight proteins, the apparent crystal density may increase with time: the crystal continues to sink and there is no apparent equilibrium spot. This behaviour is seen in both organic solvent gradients and Ficoll gradients, and the reasons for it are unclear. It may be that, in organic solvent gradients, some of the solvent can dissolve into the crystal; or the crystal may condense from slow desiccation. In Ficoll gradients, it may be that sucrose monomers or dimers are present, which diffuse into the crystal over time. A careful study of this behaviour (Bode & Schirmer, 1985) in Ficoll gradients suggested that useful density values can still be obtained for these crystals by fitting the apparent density to an exponential curve:

$$\rho_c(t) = a + b \exp(-\lambda t). \quad (5.2.6.3)$$

In this expression, parameters  $a$ ,  $b$  and  $\lambda$  must be derived from the fitted curve. The crystals were inserted into the gradient with flame-narrowed Pasteur pipettes. Each crystal was initially surrounded by a small amount of mother liquor, which rapidly diffused into the Ficoll solution. Time zero was assigned as the time when centrifugation first began. It was necessary to observe crystal positions within the first minute, and at two- to five-minute intervals thereafter, to obtain a reasonable time curve for the density function. The experimental goal in the Bode & Schirmer experiment was to obtain a good estimate for the density value at time zero,  $\rho_c(0) = a + b$ . This was realized in all six of the crystal forms that manifested time-dependent density drift in the study.

## 5.2.7. How to handle the solvent density

It is necessary to have an accurate estimate of the mean solvent density,  $\rho_s$ , in (5.2.4.9). The Ficoll gradient-tube method is particularly convenient for this reason: the gradient can be made without any significant solute other than Ficoll. Since the free-solvent compartment of the crystal is entirely water,  $\rho_s = \rho_{bs} = \rho_{fs} = 1.0 \text{ g ml}^{-1}$ . Therefore, in Ficoll density gradients, the crystal density becomes  $\rho_o$ , as defined in (5.2.5.2), and the packing number  $n$  can be calculated from

$$n = \frac{VN_o (\rho_c - 1)}{M (1 - \bar{v}_m)}. \quad (5.2.7.1)$$

Another way to set  $\rho_s = 1.0 \text{ g ml}^{-1}$  is to cross-link the crystals with glutaraldehyde (Quiocho & Richards, 1964; Cornick *et al.*, 1973; Matthews, 1985), making the crystals insoluble even in the absence of stabilizing solutes. Once cross-linked, crystals can be transferred to a water solution prior to the density measurement, thereby substituting water for its free solvent. Care must be taken with cross-linking, however. Overnight soaking in 2% glutaraldehyde solutions can substantially increase the crystal density, while destroying its crystalline order (Matthews, 1985). Even 0.5% glutaraldehyde concentrations may change the observed density of some crystals if the exposure is for many hours – which may be necessary to render the crystal completely insoluble. Therefore, the densities observed from cross-linked crystals should be regarded with caution.

If it is necessary to carry out density measurements in an organic solvent gradient, then it is necessary in general to measure the crystal density at more than one free solvent density, since the relative volume fractions of the crystal's components are not known *a priori*. However, if this is a well behaved protein crystal, by setting  $\bar{v}_m = 0.74 \text{ ml g}^{-1}$ ,  $V_M = 2.4 \text{ \AA}^3 \text{ Da}^{-1}$  and  $w = 0.25 \text{ g bound water per g protein}$ , one can guess the crystal's volume compartments to be:

$$\varphi_m = 0.51, \quad \varphi_{fs} = 0.32, \quad \varphi_{bs} = 0.17,$$

and the mean solvent density to use in (5.2.4.9) would be

$$\rho_s \approx 0.35 + 0.65\rho_{fs}. \quad (5.2.7.2)$$

This may give reasonably reliable derivations for  $n$  in (5.2.4.9), with just one crystal-density measurement. Over-reliance on parameter estimates, however, can lead to bogus results, and (5.2.7.2) should be used with caution.

## Acknowledgements

This work was supported by the US Department of Energy, Office of Biological and Environmental Research, under contract W31-109-ENG-38.

## References

- Åkervall, K. & Strandberg, B. (1971). *X-ray diffraction studies of the satellite tobacco necrosis virus. III. A new crystal mounting method allowing photographic recording of 3 Å diffraction data.* *J. Mol. Biol.* **62**, 625–627.
- Blundell, T. L. & Johnson, L. N. (1976). *Protein crystallography.* New York, London, San Francisco: Academic Press.
- Breyer, W. A., Kingston, R. L., Anderson, B. F. & Baker, E. N. (1999). *On the molecular-replacement problem in the presence of merohedral twinning: structure of the N-terminal half-molecule of human lactoferrin.* *Acta Cryst.* **D55**, 129–138.
- Britton, D. (1972). *Estimation of twinning parameter for twins with exactly superimposed reciprocal lattices.* *Acta Cryst.* **A28**, 296–297.
- Bunn, C. W. (1945). *Chemical crystallography. An introduction to optical and X-ray methods.* Oxford: Clarendon Press.
- Ducruix, A. & Geigé, R. (1992). Editors. *Crystallization of nucleic acids and proteins. A practical approach.* Oxford, New York, Tokyo: IRL Press.
- Dyda, F., Hickman, A. B., Jenkins, T. M., Engelman, A., Craigie, R. & Davies, D. R. (1994). *Crystal structure of the catalytic domain of HIV-1 integrase: similarity to other polynucleotidyl transferases.* *Science*, **266**, 1981–1986.
- Frey, M., Genovesio-Taverne, J.-C. & Fontecilla-Camps, J. C. (1988). *Application of the periodic bond chain (PBC) theory to the analysis of the molecular packing in protein crystals.* *J. Cryst. Growth*, **90**, 245–258.
- Gomis-Rüth, F. X., Fita, I., Kiefersauer, R., Huber, R., Avilés, F. X. & Navaza, J. (1995). *Determination of hemihedral twinning and initial structural analysis of crystals of the procarboxypeptidase A ternary complex.* *Acta Cryst.* **D51**, 819–823.
- Hartman, P. & Perdok, W. G. (1955a). *On the relations between structure and morphology of crystals. I.* *Acta Cryst.* **8**, 49–52.
- Hartman, P. & Perdok, W. G. (1955b). *On the relations between structure and morphology of crystals. II.* *Acta Cryst.* **8**, 521–524.
- Hartman, P. & Perdok, W. G. (1955c). *On the relations between structure and morphology of crystals. III.* *Acta Cryst.* **8**, 525–529.
- Hartshorne, N. H. & Stuart, A. (1960). *Crystals and the polarising microscope. A handbook for chemists and others*, 3rd ed. London: Edward Arnold & Co.
- Igarashi, N., Moriyama, H., Mikami, T. & Tanaka, N. (1997). *Detwinning of hemihedrally twinned crystals by the least-squares method and its application to a crystal of hydroxylamine*

## 5. CRYSTAL PROPERTIES AND HANDLING

### 5.1 (cont.)

- oxidoreductase from *Nitrosomonas europaea*. *J. Appl. Cryst.* **30**, 362–367.
- Kerfeld, C. A., Wu, Y. P., Chan, C., Krogmann, D. W. & Yeates, T. O. (1997). Crystals of the carotenoid protein from *Arthrospira maxima* containing uniformly oriented pigment molecules. *Acta Cryst.* **D53**, 720–723.
- Kudryavtsev, A. B., Mirov, S. B., DeLucas, L. J., Nicolette, C., van der Woerd, M., Bray, T. L. & Basiev, T. T. (1998). Polarized Raman spectroscopic studies of tetragonal lysozyme single crystals. *Acta Cryst.* **D54**, 1216–1229.
- McPherson, A. & Shlichta, P. (1988). Heterogeneous and epitaxial nucleation of protein crystals on mineral surfaces. *Science*, **239**, 385–387.
- McRae, D. E. (1993). *Practical protein crystallography*, pp. 21–28. San Diego, New York: Academic Press.
- Matthews, B. W. (1968). Solvent content in protein crystals. *J. Mol. Biol.* **33**, 491–497.
- Mighell, A. D., Rodgers, J. R. & Karen, V. L. (1993). Protein symmetry: metric and crystal (a precautionary note). *J. Appl. Cryst.* **26**, 68–70.
- Nadarajah, A., Li, M. & Pusey, M. L. (1997). Growth mechanism of the (110) face of tetragonal lysozyme crystals. *Acta Cryst.* **D53**, 524–534.
- Nadarajah, A. & Pusey, M. L. (1996). Growth mechanism and morphology of tetragonal lysozyme crystals. *Acta Cryst.* **D52**, 983–996.
- Oki, H., Matsuura, Y., Komatsu, H. & Chernov, A. A. (1999). Refined structure of orthorhombic lysozyme crystallized at high temperature: correlation between morphology and intermolecular contacts. *Acta Cryst.* **D55**, 114–121.
- Perutz, M. F. (1939). Absorption spectra of single crystals of haemoglobin in polarized light. *Nature (London)*, **143**, 731–733.
- Petsko, G. (1985). Flow cell construction and use. *Methods Enzymol.* **114**, 141–146.
- Phillips, F. C. (1957). *An introduction to crystallography*, 2nd ed. London, New York, Toronto: Longmans Green & Co.
- Rayment, I. (1985). Treatment and manipulation of crystals. *Methods Enzymol.* **114**, 136–140.
- Rayment, I., Johnson, J. E. & Suck, D. (1977). A method for preventing crystal slippage in macromolecular crystallography. *J. Appl. Cryst.* **10**, 365.
- Rees, D. C. (1980). The influence of twinning by merohedry on intensity statistics. *Acta Cryst.* **A36**, 578–581.
- Reichert, E. T. & Brown, A. P. (1909). *The differentiation and specificity of corresponding proteins and other vital substances in relation to biological classification and organic evolution: the crystallography of hemoglobins*. Washington DC: Carnegie Institution of Washington. (Publication No. 116.)
- Sawyer, L. & Turner, M. A. (1992). X-ray analysis. In *Crystallization of nucleic acids and proteins. A practical approach*, edited by A. Ducruix & R. Geigé, pp. 267–274. Oxford, New York, Tokyo: IRL Press.
- Stanley, E. (1972). The identification of twins from intensity statistics. *J. Appl. Cryst.* **5**, 191–194.
- Steinrauf, L. K. (1959). Preliminary X-ray data for some new crystalline forms of  $\beta$ -lactoglobulin and hen egg-white lysozyme. *Acta Cryst.* **12**, 77–79.
- Wahlstrom, E. E. (1979). *Optical crystallography*, 5th ed. New York, Chichester, Brisbane, Toronto: John Wiley.
- Yeates, T. O. (1997). Detecting and overcoming crystal twinning. *Methods Enzymol.* **276**, 344–358.
- Zhang, K. Y. J. & Eisenberg, D. (1994). Solid-state phase transition in the crystal structure of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Acta Cryst.* **D50**, 258–262.
- Berman, H. (1939). A torsion microbalance for the determination of specific gravities of minerals. *Am. Mineral.* **24**, 434–440.
- Bernal, J. D. & Crowfoot, D. (1934). Use of the centrifuge in determining the density of small crystals. *Nature (London)*, **134**, 809–810.
- Bode, W. & Schirmer, T. (1985). Determination of the protein content of crystals formed by *Mastigocladus laminosus* C-phycoerythrin, *Chroomonas spec.* phycoerythrin-645, and modified human fibrinogen using an improved Ficoll density gradient method. *Biol. Chem. Hoppe-Seyler*, **366**, 287–295.
- Cantor, C. R. & Schimmel, P. R. (1980). *Biophysical chemistry*. San Francisco: W. H. Freeman & Co.
- Coleman, P. M. & Matthews, B. W. (1971). Symmetry, molecular weight, and crystallographic data for sweet potato  $\beta$ -amylase. *J. Mol. Biol.* **60**, 163–168.
- Cornick, G., Sigler, P. B. & Ginsberg, H. S. (1973). Characterization of crystals of type 5 adenovirus hexon. *J. Mol. Biol.* **73**, 533–538.
- Crick, F. (1957). X-ray diffraction of protein crystals. *Methods Enzymol.* **4**, 127–146.
- Edelstein, S. J. & Schachman, H. (1973). Measurement of partial specific volumes by sedimentation equilibrium in  $H_2O - D_2O$  solutions. *Methods Enzymol.* **27**, 83–98.
- Edsall, J. T. (1953). Solvation of proteins. In *The proteins*, edited by H. Neurath & K. Bailey, Vol. 1, part B, pp. 549–726. New York: Academic Press.
- Graubner, H. (1986). Densitometer for absolute measurements of the temperature dependence of density, partial volumes, and thermal expansivity of solids and liquids. *Rev. Sci. Instrum.* **57**, 2817–2826.
- Hegerl, R. & Altbauer, A. (1982). The EM program system. *Ultramicroscopy*, **9**, 109–116.
- Kiefersauer, R., Stetefeld, J., Gomis-Rüth, F. X., Romão, M. J., Lottspeich, F. & Huber, R. (1996). Protein-crystal density by volume measurement and amino-acid analysis. *J. Appl. Cryst.* **29**, 311–317.
- Kratky, O., Leopold, H. & Stabinger, H. (1973). The determination of the partial specific volume of proteins by the mechanical oscillator technique. *Methods Enzymol.* **27**, 98–110.
- Kuntz, I. D. & Kaufmann, W. (1974). Hydration of proteins and polypeptides. *Adv. Protein Chem.* **28**, 239–345.
- Low, B. W. & Richards, F. M. (1952a). The use of the gradient tube for the determination of crystal densities. *J. Am. Chem. Soc.* **74**, 1660–1666.
- Low, B. W. & Richards, F. M. (1952b). Determination of protein crystal densities. *Nature (London)*, **170**, 412–415.
- Matthews, B. W. (1968). Solvent content of protein crystals. *J. Mol. Biol.* **33**, 491–497.
- Matthews, B. W. (1974). Determination of molecular weight from protein crystals. *J. Mol. Biol.* **82**, 513–526.
- Matthews, B. W. (1977). Protein crystallography. In *The proteins*, edited by H. Neurath & R. L. Hill, 403–590. New York: Academic Press.
- Matthews, B. W. (1985). Determination of protein molecular weight, hydration, and packing from crystal density. *Methods Enzymol.* **114**, 176–187.
- Midgley, H. G. (1951). A quick method of determining the density of liquid mixtures. *Acta Cryst.* **4**, 565.
- Perutz, M. F. (1946). The solvent content of protein crystals. *Trans. Faraday Soc.* **42B**, 187–195.
- Phillips, G. N., Lattman, E. E., Cummins, P., Lee, K. Y. & Cohen, C. (1979). Crystal structure and molecular interactions of tropomyosin. *Nature (London)*, **278**, 413–417.
- Quijcho, F. A. & Richards, F. M. (1964). Intermolecular cross linking of a protein in the crystalline state: carboxypeptidase-A. *Proc. Natl Acad. Sci. USA*, **52**, 833–839.
- Reilly, J. & Rae, W. N. (1954). Determination of the densities of solids by volumetry. In *Physico-chemical methods*, Vol. 1, pp. 577–608. New York: van Nostrand.
- Richards, F. M. (1954). A microbalance for the determination of protein crystal densities. *Rev. Sci. Instrum.* **24**, 1029–1034.
- Richards, F. M. & Lindley, P. F. (1999). Determination of the density of solids. In *International tables for crystallography*, Vol. C.

### 5.2

- Adair, G. S. & Adair, M. E. (1936). The densities of protein crystals and the hydration of proteins. *Proc. R. Soc. London Ser. B*, **120**, 422–446.