

3.6. CLASSIFICATION AND USE OF MACROMOLECULAR DATA

PHASING_MAD_EXPT
 PHASING_MAD_RATIO
 PHASING_MAD_SET

Phasing via multiple isomorphous replacement (§3.6.6.1.5)

PHASING_MIR
 PHASING_MIR_DER
 PHASING_MIR_DER_REFLN
 PHASING_MIR_DER_SHELL
 PHASING_MIR_DER_SITE
 PHASING_MIR_DER_SHELL

Phasing data sets (§3.6.6.1.6)

PHASING_SET
 PHASING_SET_REFLN

The data items in the PHASING category group can be used to record details about the phasing of the structure and cover the various methods used in the phasing process. Many data items are provided for multiple isomorphous replacement (MIR) and multiple-wavelength anomalous dispersion (MAD). More limited sets of data items are provided for phasing using molecular averaging and phasing *via* using a structure that is isomorphous to the present structure. The current version of the mmCIF dictionary does not provide specific data items for recording the details of phasing *via* molecular replacement.

3.6.6.1.1. Overall description of phasing

The single data item in this category is as follows:

PHASING
 • `_phasing.method`

The bullet (•) indicates a category key.

Phasing of macromolecular structures often involves the application of more than one of the methods described in the PHASING section of the mmCIF dictionary, such as when phases generated from a multiple isomorphous replacement experiment are improved by molecular averaging. The PHASING category is used to list the methods that were used.

At present, the category contains a single data item, the purpose of which is to specify the method employed in the structure determination. It may have one or more of the values listed in the dictionary (Example 3.6.6.1).

3.6.6.1.2. Phasing via molecular averaging

The data items in this category are as follows:

PHASING_AVERAGING
 • `_phasing_averaging.entry_id`
 → `_entry.id`
`_phasing_averaging.details`
`_phasing_averaging.method`

The bullet (•) indicates a category key. The arrow (→) is a reference to a parent data item.

When more than one copy of a molecule is present in the asymmetric unit, phases can be improved by averaging an electron-density map over the multiple images of the molecule. In some special cases with very high noncrystallographic symmetry, *de novo* phases have been derived by iterative application of molecular averaging, but more often averaging is used to improve phases determined by another method.

There are many protocols used for phasing with averaging and they are very varied. It was not thought to be appropriate to specify data items for any one approach in the current version of the mmCIF dictionary. The data items that are provided allow a text-based description of the protocol to be given; a formalism

Example 3.6.6.1. The methods used to generate the phases for a hypothetical structure described with the data item in the PHASING category.

```
loop_
  _phasing.method
    'mir'
    'averaging'
```

Example 3.6.6.2. Phase improvement with molecular averaging for a hypothetical structure described with data items in the PHASING_AVERAGING category.

```
_phasing_averaging.entry_id    'EXAMHYPO'
_phasing_averaging.method
; Iterative threefold averaging alternating with
  phase extensions by 0.5 reciprocal lattice units
  per cycle.
;
_phasing_averaging.details
; The position of the threefold axis was redetermined
  every five cycles.
;
```

for recording a fully parsable description of molecular averaging needs to be developed for future revisions of the dictionary.

Data items in the PHASING_AVERAGING category allow free-text descriptions to be given of the method used for structure determination or phase improvement using averaging over multiple observations of the molecule in the asymmetric unit and of any specific details of the application of the method to the current structure determination (Example 3.6.6.2). Note that the reference to the method is to be used to describe the method itself, and not as a reference to a software package; references to software packages would be made using data items in the SOFTWARE category.

3.6.6.1.3. Phasing via isomorphous replacement

The data items in this category are as follows:

PHASING_ISOMORPHOUS
 • `_phasing_isomorphous.entry_id`
 → `_entry.id`
`_phasing_isomorphous.details`
`_phasing_isomorphous.method`
`_phasing_isomorphous.parent`

The bullet (•) indicates a category key. The arrow (→) is a reference to a parent data item.

Phases for many macromolecular structures are obtained from a previous determination of the same structure in the same crystal lattice. Examples of this are the determination of the structure of a point mutant or the determination of a structure in which a ligand is bound to an active site that was empty in the previous structure determination. In these cases, the new structure is essentially isomorphous with the parent structure, hence this method of phasing is termed 'isomorphous phasing' in the mmCIF dictionary. It is not to be confused with multiple isomorphous phasing (MIR), a phasing technique that involves the use of heavy-atom derivatives. MIR phasing is discussed in Section 3.6.6.1.5.

Not much information is needed to characterize isomorphous phasing. The 'parent' structure (the structure used to generate the initial phases for the present structure) is described in a free-text field and a second free-text field can be used to give details of the application of the method to the determination of the present structure (for instance, the removal of solvent or a bound ligand). In Example 3.6.6.3, the parent structure is the PDB entry 5HVP and the structure that is the subject of the present data block is identified as 'HVP+CmpdA'. `_phasing_isomorphous.method` allows